arXiv:1006.0464v1 [math.RA] 2 Jun 2010

CHEVALLEY SUPERGROUPS
OF TYPE $D(2,1;a)$

F. Gavarini

Dipartimento di Matematica, Università di Roma "Tor Vergata"
via della ricerca scientifica 1 — I-00133 Roma, Italy
e-mail: gavarini@mat.uniroma2.it

Abstract

We present a construction "à la Chevalley" of affine supergroups associated to Lie superalgebras of type $D(2,1;a)$, for any possible value of the parameter a. This extends the similar work performed in [4] which associates an affine supergroup to any other simple Lie superalgebras of classical type.

1 Introduction

[1] In his work of 1955, Chevalley provided a combinatorial construction of all simple affine algebraic groups over any field. In particular, his method led to an existence theorem for simple affine algebraic groups: one starts with a simple (complex, finite-dimensional) Lie algebra and a simple module V for it, and realizes the required group as a closed subgroup of $\mathrm{GL}(V)$. Furthermore, Chevalley's construction can be recast as to provide a description of all simple affine groups as group schemes over $\mathbb{Z}$.

In [4] the philosophy of Chevalley was revisited in the context of supergeometry. The outcome is a construction of affine supergroups whose tangent Lie superalgebra is of *classical type*. However, some exceptions were left out, namely the cases when the Lie superalgebra is of type $D(2,1;a)$ and the parameter a is not an integer number. The present work fills in this gap.

---

[1] 2010 *MSC*: Primary 14M30, 14A22; Secondary 17B20.

By "affine supergroup" here I mean a representable functor from the category (salg) of commutative superalgebras — over some fixed ground ring — to the category (groups) of groups: in other words, an affine supergroup-scheme, identified with its functor of points. Thus we adopt the functorial language (à la Grothendieck, say) from scratch. The general procedure developed in [4] proceeds in two steps: first, one constructs a functor from (salg) to (groups), recovering Chevalley's ideas to define the values of such a group functor on each superalgebra A — i.e., to define its A–points; second, one proves that the sheafification of this functor is representable.

For the case $D(2,1;a)$ — with $a \notin \mathbb{Z}$ — one needs a careful modification of the general procedure of [4]; thus the presentation hereafter will detail those steps which need changes, and will simply refer to [4] for those where the original arguments still work unchanged.

The initial datum is a simple Lie superalgebra $\mathfrak{g} = D(2,1;a)$.

We start proving some basic results on $\mathfrak{g}$, namely the existence of *Chevalley bases* (i.e. bases having nice integrality properties) and a PBW-like theorem for the Kostant $\mathbb{Z}$–form of the universal enveloping superalgebra $U(\mathfrak{g})$.

Next we take a faithful, finite-dimensional $\mathfrak{g}$–module V, and we show it has suitable lattices M invariant by the Kostant superalgebra. This allows to define — functorially — additive and multiplicative one-parameter (super)subgroups of operators acting on scalar extensions of M. The additive subgroups are just like in the general case: there exists one of them for every root of $\mathfrak{g}$. The multiplicative ones instead are associated to elements of the fixed Cartan subalgebra of $\mathfrak{g}$, and are of two types: those of *classical* type, modeled on the group functor $A \mapsto U(A_0)$ — the *group of units* of A_0 — and those of *a–type*, modeled on the group functor $A \mapsto P_a(A)$ — the group of elements of A_0 "which may be raised to the a^k–th power, for all k". The second type of multiplicative one-parameter subgroups were not used in [4]: here one needs them because one has to consider the power operation $t \mapsto t^a$, which in general is not defined for any $t \in U(A_0)$, but it is for $t \in P_a(A)$; this marks a difference with the integral case (i.e. for $a \in \mathbb{Z}$).

Then we consider the functor $G : (\text{salg}) \longrightarrow (\text{groups})$ whose value $G(A)$ on $A \in (\text{salg})$ is the subgroup of $\mathrm{GL}(V(A))$ — with $V(A) := A \otimes M$ — generated by all the homogeneous one-parameter supersubgroups mentioned above. This group functor is in fact a presheaf, hence we can take its sheafification $\mathbf{G}_V = \mathbf{G} : (\text{salg}) \longrightarrow (\text{groups})$ (in the sense of category theory). These $\mathbf{G}_V$ are, by definition, our "Chevalley supergroups".

Acting just like in [4], one defines a "classical affine subgroup" $\mathbf{G}_0$ of $\mathbf{G}_V$, corresponding to the even part $\mathfrak{g}_0$ of $\mathfrak{g}$ (and to V), and then finds a factorization $\mathbf{G}_V = \mathbf{G}_0 \, \mathbf{G}_1 \cong \mathbf{G}_0 \times \mathbf{G}_1$, where $\mathbf{G}_1$ corresponds instead to the odd part $\mathfrak{g}_1$ of $\mathfrak{g}$. Actually, one has even a finer factorization $\mathbf{G}_V = \mathbf{G}_0 \times \mathbf{G}_1^{-,<} \times \mathbf{G}_1^{+,<}$ with $\mathbf{G}_1^{\pm,<}$ being totally odd superspaces associated to the positive or negative odd roots of $\mathfrak{g}$. Thus $\mathbf{G}_1 = \mathbf{G}_1^{-,<} \times \mathbf{G}_1^{+,<}$ is representable, and $\mathbf{G}_0$ is representable too, hence the above factorization implies that $\mathbf{G}_V$ is representable too, so it is an affine supergroup. The outcome then is that our Chevalley supergroups are affine supergroups.

Finally, one also proves that our construction is functorial in V and that $\text{Lie}(\mathbf{G}_V)$ is just $\mathfrak{g}$ as one expects, like in [4] (no special changes are needed).

Acknowledgements

The author thanks M. Duflo, R. Fioresi and C. Gasbarri for many useful conversations, and K. Iohara and Y. Koga for their valuable explanations.

2 Preliminaries

In this section we introduce some basic preliminaries of supergeometry. For further details, our main reference is [3].

2.1 Superalgebras, superspaces, supergroups

Let $\Bbbk$ be a unital, commutative ring.

We call $\Bbbk$–*superalgebra* any associative, unital $\Bbbk$–algebra A which is $\mathbb{Z}_2$–graded; that is, A is a $\Bbbk$–algebra graded by the two-element group $\mathbb{Z}_2$. Thus A splits as $A = A_0 \oplus A_1$, and $A_a \, A_b \subseteq A_{a+b}$. The $\Bbbk$–submodule A_0 and its elements are called *even*, while A_1 and its elements *odd*. By $p(x)$ we denote the *parity* of any homogeneous element $x \in A_{p(x)}$. Clearly, $\Bbbk$–superalgebras form a category, whose morphisms are all those in the category of algebras which preserve the unit and the $\mathbb{Z}_2$–grading. At last, for any $n \in \mathbb{N}$ we call A_1^n the A_0–submodule of A spanned by all products $\vartheta_1 \cdots \vartheta_n$ with $\vartheta_i \in A_1$ for all i, and $A_1^{(n)}$ the unital subalgebra of A generated by A_1^n.

A superalgebra A is said to be *commutative* iff $xy = (-1)^{p(x)p(y)} yx$ for all homogeneous x, $y \in A$. We denote by (salg) the category of commutative superalgebras; if necessary, we shall stress the role of $\Bbbk$ by writing $(\mathrm{salg})_{\Bbbk}$.

Definition 2.1. A *superspace* $S = \big(|S|, \mathcal{O}_S\big)$ is a topological space $|S|$ with a sheaf of commutative superalgebras $\mathcal{O}_S$ such that the stalk $\mathcal{O}_{S,x}$ is a local superalgebra for all $x \in |S|$. A *morphism* $\phi : S \longrightarrow T$ of superspaces is a pair $\phi = \big(|\phi|, \phi^*\big)$, where $\phi : |S| \longrightarrow |T|$ is a morphism of topological spaces and $\phi^* : \mathcal{O}_T \longrightarrow \phi_* \mathcal{O}_S$ is a sheaf morphism such that $\phi_x^*\big(\mathbf{m}_{|\phi|(x)}\big) = \mathbf{m}_x$, where $\mathbf{m}_{|\phi|(x)}$ and $\mathbf{m}_x$ are the maximal ideals in the stalks $\mathcal{O}_{T,\,|\phi|(x)}$ and $\mathcal{O}_{S,x}$ respectively and ϕ_x^* is the morphism induced by ϕ^* on the stalks. Here as usual $\phi_* \mathcal{O}_S$ is the sheaf on $|T|$ defined as $\phi_* \mathcal{O}_S(V) := \mathcal{O}_S(\phi^{-1}(V))$.

Given a superspace $S = \big(|S|, \mathcal{O}_S\big)$, let $\mathcal{O}_{S,0}$ and $\mathcal{O}_{S,1}$ be the sheaves on $|S|$ defined as follows: $\mathcal{O}_{S,0}(U) := \mathcal{O}_S(U)_0$, $\mathcal{O}_{S,1}(U) := \mathcal{O}_S(U)_1$ for each open subset U in $|S|$. Then $\mathcal{O}_{S,0}$ is a sheaf of ordinary commutative algebras, while $\mathcal{O}_{S,1}$ is a sheaf of $\mathcal{O}_{S,0}$–modules.

Definition 2.2. A *superscheme* is a superspace $S := \big(|S|, \mathcal{O}_S\big)$ such that $\big(|S|, \mathcal{O}_{S,0}\big)$ is an ordinary scheme and $\mathcal{O}_{S,1}$ is a quasi-coherent sheaf of $\mathcal{O}_{S,0}$–modules. A *morphism* of superschemes is a morphism of the underlying superspaces.

Definition 2.3. Let $A \in (\mathrm{salg})$ and let $\mathcal{O}_{A_0}$ be the structural sheaf of the ordinary scheme $\underline{\mathit{Spec}}\,(A_0) = \big(\mathit{Spec}\,(A_0), \mathcal{O}_{A_0}\big)$, where $\mathit{Spec}\,(A_0)$ denotes the prime spectrum of the commutative ring A_0 . Now A is a module over A_0 , so we have a sheaf $\mathcal{O}_A$ of $\mathcal{O}_{A_0}$–modules over $\mathit{Spec}\,(A_0)$ with stalk A_p , the p–localization of the A_0–module A , at the prime $p \in \mathit{Spec}\,(A_0)$.

We define the superspace $\underline{\mathit{Spec}}\,(A) := \big(\mathit{Spec}\,(A_0), \mathcal{O}_A\big)$. By definition, $\underline{\mathit{Spec}}\,(A)$ is a superscheme; we call *affine* any superscheme which is isomorphic to $\underline{\mathit{Spec}}\,(A)$ for some commutative superalgebra A .

Clearly any superscheme is locally isomorphic to an affine superscheme.

Example 2.4. The *affine superspace* $\mathbb{A}_{\Bbbk}^{p|q}$, also denoted $\Bbbk^{p|q}$, is defined — for each p , $q \in \mathbb{N}$ — as $\mathbb{A}_{\Bbbk}^{p|q} := \underline{\mathit{Spec}}\big(\Bbbk[x_1, \dots, x_p] \otimes_{\Bbbk} \Bbbk[\xi_1 \dots \xi_q]\big)$, where $\Bbbk[\xi_1 \dots \xi_q]$ is the exterior (or "Grassmann") algebra generated by $\xi_1, \dots, \xi_q$, and $\Bbbk[x_1, \dots, x_p]$ the polynomial algebra in p commuting indeterminates. $\diamondsuit$

Definition 2.5. Let X be a superscheme. Its *functor of points* is the functor $h_X : (\text{salg}) \longrightarrow (\text{sets})$ defined on objects by $h_X(A) := \text{Hom}\big(\underline{Spec}(A), X\big)$ and on arrows by $h_X(f)(\phi) := \phi \circ \underline{Spec}(f)$. If (groups) is the category of groups and h_X is a functor $h_X : (\text{salg}) \longrightarrow (\text{groups})$, then we say that X is a *supergroup*. When X is affine, this is equivalent to the fact that $\mathcal{O}(X)$ — the superalgebra of global sections of the structure sheaf on X — is a (commutative) *Hopf superalgebra*. More in general, by *supergroup functor* we mean any functor $G : (\text{salg}) \longrightarrow (\text{groups})$.

Any representable supergroup functor is the same as an affine supergroup: indeed, the former corresponds to the functor of points of the latter. Indeed, the functor of points of a given superscheme recaptures all the information carried by a superscheme (by Yoneda's Lemma): in particular, two superschemes are isomorphic if and only if their functors of points are. Thus we use the same letter to denote both a superscheme and its functor of points.

More details can be found in [3], Ch. 3–5.

In the present work, we shall actually consider only affine supergroups, which we are going to describe via their functor of points.

The next examples turn out to be very important in the sequel.

Examples 2.6.

(1) Let V be a super vector space. For any superalgebra A we define $V(A) := (A \otimes V)_0 = A_0 \otimes V_0 \oplus A_1 \otimes V_1$. This is a representable functor in the category of superalgebras, whose representing object is the superalgebra of polynomial functions on V . Hence V can be seen as an affine superscheme.

(2) $\text{GL}(V)$ *as an affine supergroup.* Let V be a finite dimensional super vector space of dimension $p|q$. For any superalgebra A , let $\text{GL}(V)(A) := \text{GL}\big(V(A)\big)$ be the set of isomorphisms $V(A) \longrightarrow V(A)$. If we fix a homogeneous basis for V, we see that $V \cong \Bbbk^{p|q}$; in other words, $V_0 \cong \Bbbk^p$ and $V_1 \cong \Bbbk^q$. In this case, we also denote $\text{GL}(V)$ with $\text{GL}(p|q)$. Now, $\text{GL}(p|q)(A)$ is the group of invertible matrices of size $(p+q)$ with diagonal block entries in A_0 and off-diagonal block entries in A_1 . It is known that the functor $\text{GL}(V)$ is representable; see (e.g.), [14], Ch. 3, for further details. $\diamondsuit$

2.2 Lie superalgebras

From now on, we assume $\Bbbk$ to be such that 2 and 3 are not zero and they are not zero divisors in $\Bbbk$. Moreover, *all $\Bbbk$–modules hereafter will be assumed to have no p–torsion, for* $p \in \{2,3\}$.

Definition 2.7. Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be a super (i.e., $\mathbb{Z}_2$–graded) $\Bbbk$–module (with no p–torsion for $p \in \{2,3\}$, as mentioned above). We say that $\mathfrak{g}$ is a Lie superalgebra, if we have a bracket $[\ ,\ ] : \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g}$ which satisfies the following properties (for all $x, y \in \mathfrak{g}$ homogeneous):

(1) Anti-symmetry: $\qquad [x,y] + (-1)^{p(x)\,p(y)}[y,x] = 0$

(2) Jacobi identity:

$$(-1)^{p(x)\,p(z)}\,[x,[y,z]] + (-1)^{p(y)\,p(x)}\,[y,[z,x]] + (-1)^{p(z)\,p(y)}\,[z,[x,y]] = 0$$

Example 2.8. Let $V = V_0 \oplus V_1$ be a *free* super $\Bbbk$–module, and consider $\mathrm{End}(V)$, the endomorphisms of V as an ordinary $\Bbbk$–module. This is again a free super $\Bbbk$–module, $\mathrm{End}(V) = \mathrm{End}(V)_0 \oplus \mathrm{End}(V)_1$, where $\mathrm{End}(V)_0$ are the morphisms which preserve the parity, while $\mathrm{End}(V)_1$ are the morphisms which reverse the parity. If V has finite rank, and we choose a basis for V of homogeneous elements (writing first the even ones), then $\mathrm{End}(V)_0$ is the set of all diagonal block matrices, while $\mathrm{End}(V)_1$ is the set of all off-diagonal block matrices. Thus $\mathrm{End}(V)$ is a Lie superalgebra with bracket

$$[A,B] := AB - (-1)^{|A||B|}\,BA \qquad \text{for all homogeneous } A, B \in \mathrm{End}(V)\,.$$

The standard example is $V := \Bbbk^{p|q} = \Bbbk^p \oplus \Bbbk^q$, with $V_0 := \Bbbk^p$ and $V_1 := \Bbbk^q$. In this case we also write $\mathrm{End}\big(\Bbbk^{m|n}\big) := \mathrm{End}(V)$ or $\mathfrak{gl}(p\,|q) := \mathrm{End}(V)$. ◊

A Lie superalgebra $\mathfrak{g}$ is called *classical* if it is simple, i.e. it has no non-trivial (homogeneous) ideals, and $\mathfrak{g}_1$ is semisimple as a $\mathfrak{g}_0$–module. Classical Lie superalgebras of finite dimension over algebraically closed fields of characteristic zero were classified by V. Kac (cf. [8], [10]).

We assume the reader is familiar with the standard, basic terminology and notions about classical Lie superalgebras ("Cartan subalgebra(s)", "roots", "coroots", "root vectors", etc). The main reference for that is [8], but for the purposes of the present work one can also refer to [4].

2.3 The Lie superalgebra $D(2,1;a)$

Let $\mathbb{K}$ be an algebraically closed field of characteristic zero, and let $a \in \mathbb{K} \setminus \{0,-1\}$. Then let $\mathbb{Z}[a]$ be the unital subring of $\mathbb{K}$ generated by a. Clearly $\mathbb{Z}[a] = \mathbb{Z}$ if and only if $a \in \mathbb{Z}$.

According to Kac's work, we can realize $\mathfrak{g} := D(2,1;a)$ as a contragredient Lie superalgebra: in particular, it admits a presentation by generators and relations with a standard procedure (detailed in general in [8]). In order to do that, we first fix a specific choice of Dynkin diagram and corresponding Cartan matrix, like in [5], §2.28 (first choice), namely



$$, \qquad \left(a_{i,j}\right)_{i,j=1,2,3;} := \begin{pmatrix} 0 & 1 & a \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$$

Now we define $\mathfrak{g} = D(2,1;a)$ as the Lie superalgebra over $\mathbb{K}$ with generators h_i, e_i, f_i $(i = 1,2,3)$, with degrees $p(h_i) := 0$, $p(e_i) := \delta_{1,i}$, $p(f_i) := \delta_{1,i}$ $(i = 1,2,3)$, and with relations (for all $i, j = 1,2,3$)

$$\begin{gathered} \big[h_i, h_j\big] = 0 \ , \qquad \big[e_1, e_1\big] = 0 \ , \qquad \big[f_1, f_1\big] = 0 \ , \\ \big[h_i, e_j\big] = +a_{i,j}\, e_j \ , \qquad \big[h_i, f_j\big] = -a_{i,j}\, f_j \ , \qquad \big[e_i, f_j\big] = \delta_{i,j}\, h_i \ . \end{gathered}$$

The subspace $\mathfrak{h} := \sum_{i=1}^{3} \mathbb{K}\, h_i$ is a *Cartan subalgebra* of $\mathfrak{g}$ (included in $\mathfrak{g}_0$). Then the adjoint action of $\mathfrak{h}$ splits $\mathfrak{g}$ into eigenspaces, namely

$$\mathfrak{g} = \bigoplus_{\alpha\in\mathfrak{h}^*} \mathfrak{g}_\alpha \ , \qquad \text{with} \quad \mathfrak{g}_\alpha := \big\{ x \in \mathfrak{g} \,\big|\, [h,x] = \alpha(h)\, x \,,\ \forall\, h \in \mathfrak{h} \big\} \quad \forall\, \alpha \in \mathfrak{h}^*$$

Then we define the *roots* of $\mathfrak{g}$ by $\Delta := \Delta_0 \coprod \Delta_1 = \{\, \textit{roots} \text{ of } \mathfrak{g} \,\}$ with

$$\begin{gathered} \Delta_0 := \big\{\, \alpha \in \mathfrak{h}^* \setminus \{0\} \,\big|\, \mathfrak{g}_\alpha \cap \mathfrak{g}_0 \neq \{0\} \big\} = \{\, \textit{even roots} \text{ of } \mathfrak{g} \,\} \\ \Delta_1 := \big\{\, \alpha \in \mathfrak{h}^* \,\big|\, \mathfrak{g}_\alpha \cap \mathfrak{g}_1 \neq \{0\} \big\} = \{\, \textit{odd roots} \text{ of } \mathfrak{g} \,\} \end{gathered}$$

Δ is called the *root system* of $\mathfrak{g}$, and for each root α we call $\mathfrak{g}_\alpha$ its *root space.* Moreover, every non-zero vector in a root space is called *root vector.*

In the present case, every root space is one dimensional, so any root vector forms a basis of its own root space. Then any $\mathbb{K}$–basis of $\mathfrak{h}$ together with any choice of a root vector for each root will provide a $\mathbb{K}$–basis of $\mathfrak{g} = D(2,1;a)$.

There is an even, non-degenerate, invariant bilinear form on $\mathfrak{g}$, whose restriction to $\mathfrak{h}$ is in turn an invariant bilinear form on $\mathfrak{h}$. We denote this form by $\big(x,y\big)$, and we use it to identify $\mathfrak{h}^*$ to $\mathfrak{h}$, via $\alpha \mapsto H_\alpha$, and then to define a similar form on $\mathfrak{h}^*$, such that $\big(\alpha',\alpha''\big) = \big(H_{\alpha'},H_{\alpha''}\big)$. In particular, we fix normalizations so that $\alpha(H_\alpha)=2$ and $\alpha(H_{\alpha'})=0$ for all $\alpha,\alpha'\in\Delta_0$ such that $\big(\alpha\,,\alpha'\big)=0$ (in short, we adopt the normalizations as in [6]). Moreover, if α is any (even, odd, positive or negative) root we shall call the vector H_α the (even, odd, positive or negative) *coroot* associated to α .

Actually, we can describe explicitly the root system of $\mathfrak{g} = D(2,1;a)$ (after [5], Table 3.60) as follows:

$$\Delta \;=\; \big\{\, \pm 2\,\varepsilon_1 \,,\, \pm 2\,\varepsilon_2 \,,\, \pm 2\,\varepsilon_3 \,,\, \pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \,\big\}$$

$$\Delta_0 \;=\; \big\{\, \pm 2\,\varepsilon_1 \,,\, \pm 2\,\varepsilon_2 \,,\, \pm 2\,\varepsilon_3 \,\big\} \;\;, \qquad \Delta_1 \;=\; \big\{\, \pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \,\big\}$$

where $\{\varepsilon_1,\varepsilon_2,\varepsilon_3\}$ is an orthogonal basis in a $\mathbb{K}$–vector space with inner product $(\ ,\ )$ such that $(\varepsilon_1,\varepsilon_1) = -(1+a)\big/2$, $(\varepsilon_2,\varepsilon_2) = 1\big/2$, $(\varepsilon_3,\varepsilon_3) = a\big/2$.

According to this choice, we can fix a distinguished system of *simple roots*, say $\{\alpha_1,\alpha_2,\alpha_3\}$, associated to it, namely

$$\alpha_1 \;=\; \varepsilon_1 - \varepsilon_2 - \varepsilon_3 \;\;, \quad \alpha_2 \;=\; 2\,\varepsilon_2 \;\;, \quad \alpha_3 \;=\; 2\,\varepsilon_3 \;\;.$$

In terms of these, we call *positive* the roots

$$2\,\varepsilon_1 \;=\; 2\,\alpha_1 + \alpha_2 + \alpha_3 \;\;, \qquad 2\,\varepsilon_2 \;=\; \alpha_2 \;\;, \qquad 2\,\varepsilon_3 \;=\; \alpha_3$$

$$\varepsilon_1 - \varepsilon_2 - \varepsilon_3 \;=\; \alpha_1 \;\;, \qquad \varepsilon_1 + \varepsilon_2 - \varepsilon_3 \;=\; \alpha_1 + \alpha_2 \;\;,$$

$$\varepsilon_1 - \varepsilon_2 + \varepsilon_3 \;=\; \alpha_1 + \alpha_3 \;\;, \qquad \varepsilon_1 + \varepsilon_2 + \varepsilon_3 \;=\; \alpha_1 + \alpha_2 + \alpha_3$$

(those in first line being even, the others odd), calling their set by Δ^+ : so

$$\Delta^+ \;=\; \{\alpha_1 \,,\, \alpha_2 \,,\, \alpha_3 \,,\, \alpha_1+\alpha_2 \,,\, \alpha_1+\alpha_3 \,,\, \alpha_1+\alpha_2+\alpha_3 \,,\, 2\,\alpha_1+\alpha_2+\alpha_3\}$$

We call instead *negative* the roots in $\Delta^- := -\Delta^+$. So the root system is given by $\Delta = \Delta^+ \coprod \Delta^-$. We also set $\Delta_0^\pm := \Delta_0 \cap \Delta^\pm$, $\Delta_1^\pm := \Delta_1 \cap \Delta^\pm$.

It is worth stressing at this point that the coroots $H_\alpha \in \mathfrak{h}$ associated to the positive roots are $H_{2\varepsilon_1} = (1+a)^{-1}\big(2\,h_1 - h_2 - a\,h_3\big)$, $H_{2\varepsilon_2} = h_2$, $H_{2\varepsilon_3} = h_3$, $H_{\varepsilon_1-\varepsilon_2-\varepsilon_3} = h_1$, $H_{\varepsilon_1+\varepsilon_2-\varepsilon_3} = h_1 - h_2$, $H_{\varepsilon_1-\varepsilon_2+\varepsilon_3} = h_1 - a\,h_3$, and $H_{\varepsilon_1+\varepsilon_2+\varepsilon_3} = h_1 - h_2 - a\,h_3$; then the formula $H_{-\alpha} = H_\alpha$ yields coroots associated to negative roots out of those associated to positive ones.

Now we introduce the following elements:

$$e_{1,2} := \big[e_1,e_2\big] \;,\quad e_{1,3} := \big[e_1,e_3\big] \;,\quad e_{1,2,3} := \big[e_{1,2},e_3\big] \;,\quad e'_{1,1,2,3} := \big[e_1,e_{1,2,3}\big]$$

$$f_{2,1} := \big[f_2,f_1\big] \;,\quad f_{3,1} := \big[f_3,f_1\big] \;,\quad e_{3,2,1} := \big[f_3,f_{2,1}\big] \;,\quad f'_{3,2,1,1} := \big[f_{3,2,1},f_1\big]$$

All these are root vectors, respectively for the positive roots $\alpha_1+\alpha_2$, $\alpha_1+\alpha_3$, $\alpha_1+\alpha_2+\alpha_3$ and $2\,\alpha_1+\alpha_2+\alpha_3$ (for the first line vectors) and similarly for the negative roots (for the second line vectors). Moreover, by definition the generators e_i and f_i ($i=1,2,3$) are root vectors respectively for the roots $+\alpha_i$ and $-\alpha_i$ ($i=1,2,3$). As $\{h_1,h_2,h_3\}$ is a $\mathbb{K}$–basis of $\mathfrak{h}$, we conclude that all these root vectors along with h_1 , h_2 and h_3 form a $\mathbb{K}$–basis of $\mathfrak{g}$.

The relevant new brackets among all these basis elements — dropping the zero ones, those given by the very definition of $D(2,1;a)$, those coming from others by (super-)skewcommutativity, and those involving the h_i's (which are given by the fact that all involved vectors are eigenvectors for $\mathfrak{h}$ — are

$$\big[e_1,e_2\big]=e_{1,2}\ ,\quad \big[e_1,e_3\big]=e_{1,3}\ ,\quad \big[e_1,e_{1,2,3}\big]=e'_{1,1,2,3}$$

$$\big[e_1,f_{2,1}\big]=f_2\ ,\quad \big[e_1,f_{3,1}\big]=af_3\ ,\quad \big[e_1,f'_{3,2,1,1}\big]=-(1+a)f_{3,2,1}$$

$$\big[e_2,e_{1,3}\big]=-e_{1,2,3}\ ,\quad \big[e_2,f_{2,1}\big]=f_1\ ,\quad \big[e_2,f_{3,2,1}\big]=f_{3,1}$$

$$\big[e_3,e_{1,2}\big]=-e_{1,2,3}\ ,\quad \big[e_3,f_{3,1}\big]=f_1\ ,\quad \big[e_3,f_{3,2,1}\big]=f_{2,1}$$

$$\big[f_1,f_2\big]=-f_{2,1}\ ,\quad \big[f_1,f_3\big]=-f_{3,1}\ ,\quad \big[f_1,f_{3,2,1}\big]=f'_{3,2,1,1}$$

$$\big[f_1,e_{1,2}\big]=e_2\ ,\quad \big[f_1,e_{1,3}\big]=a\,e_3\ ,\quad \big[f_1,e'_{1,1,2,3}\big]=(1+a)\,e_{1,2,3}$$

$$\big[f_2,f_{3,1}\big]=f_{3,2,1}\ ,\quad \big[f_2,e_{1,2}\big]=-e_1\ ,\quad \big[f_2,e_{1,2,3}\big]=-e_{1,3}$$

$$\big[f_3,f_{2,1}\big]=f_{3,2,1}\ ,\quad \big[f_3,e_{1,3}\big]=-e_1\ ,\quad \big[f_3,e_{1,2,3}\big]=-e_{1,2}$$

$$\big[e_{1,2},e_{1,3}\big]=-e'_{1,1,2,3}\ ,\qquad \big[e_{1,2},f_{2,1}\big]\ =\ h_1-h_2\ ,$$

$$\big[e_{1,2},f_{3,2,1}\big]=af_3\ ,\qquad \big[e_{1,2},f'_{3,2,1,1}\big]=(1+a)f_{3,1}$$

$$\big[e_{1,3},f_{3,1}\big]\ =\ h_1-ah_3\ ,\quad \big[e_{1,3},f_{3,2,1}\big]=f_2\ ,\quad \big[e_{1,3},f'_{3,2,1,1}\big]=(1+a)f_{2,1}$$

$$\big[f_{2,1},f_{3,1}\big]=-f'_{3,2,1,1}\ ,\quad \big[f_{2,1},e_{1,2,3}\big]=a\,e_3\ ,\quad \big[f_{2,1},e'_{1,1,2,3}\big]=-(1+a)\,e_{1,3}$$

$$\big[f_{3,1},e_{1,2,3}\big]=e_2\ ,\quad \big[f_{3,1},e'_{1,1,2,3}\big]=-(1+a)\,e_{1,2}$$

$$\big[e_{1,2,3},f_{3,2,1}\big]=h_1-h_2-ah_3\ ,\quad \big[e_{1,2,3},f'_{3,2,1,1}\big]=-(1+a)f_1\ ,$$

$$\big[f_{3,2,1},e'_{1,1,2,3}\big]=-(1+a)\,e_1\ ,\quad \big[e'_{1,1,2,3},f'_{3,2,1,1}\big]=-(1+a)\big(2\,h_1-h_2-ah_3\big)$$

Now we modify just two root vectors taking

$$e_{1,1,2,3}\ :=\ +(1+a)^{-1}\,e'_{1,1,2,3}\quad ,\qquad f_{3,2,1,1}\ :=\ -(1+a)^{-1}f'_{3,2,1,1}\quad ;$$

(recall that $a \neq -1$ by assumption); then the above formulas has to be modified accordingly (plenty of coefficients $(1+a)$ just cancel out).

In particular, one can now check that the even part of $\mathfrak{g} := D(2,1;a)$, is $\mathfrak{g}_0 = \mathfrak{sl}_2 \oplus \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$. Moreover, the three triples

$$\big(e_{1,1,2,3}\,,\, f_{3,2,1,1}\,,\, (1+a)^{-1}(2\,h_1 - h_2 - a\,h_3)\big)\;, \quad \big(e_2\,,\, f_2\,,\, h_2\big)\;, \quad \big(e_3\,,\, f_3\,,\, h_3\big)$$

are $\mathfrak{sl}_2$–triples inside $\mathfrak{g}$, each one being associated to a (positive) even root $2\,\varepsilon_i$ ($i = 1,2,3$) and spanning a copy of $\mathfrak{sl}_2$ inside $\mathfrak{g}_0$.

3 Chevalley bases and Kostant superalgebras for $D(2,1;a)$

In this section we introduce the first results we shall build upon to construct our Chevalley supergroups of type $D(2,1;a)$. As before, $\mathbb{K}$ will be an algebraically closed field of characteristic zero.

3.1 Chevalley bases and Chevalley Lie superalgebras

The subject of this subsection is an analogue, in the super setting, of a classical result due to Chevalley: the notion of Chevalley basis, and corrispondingly of Chevalley Lie superalgebra. For $\mathfrak{g} := D(2,1;a)$, this notion is introduced exactly like in Definition 3.3 in [4], *up to changing* $\mathbb{Z}$ *to* $\mathbb{Z}[a]$, the latter being the unital subring of $\mathbb{K}$ generated by a (cf. §2.3).

Definition 3.1. We call *Chevalley basis* of $\mathfrak{g} := D(2,1;a)$ any homogeneous $\mathbb{K}$–basis $B = \big\{H_i\big\}_{1,2,3} \coprod \big\{X_\alpha\big\}_{\alpha\in\Delta}$ of $\mathfrak{g}$ with the following properties:

(a) $\big\{H_1, H_2, H_3\big\}$ is a $\mathbb{K}$–basis of $\mathfrak{h}$; moreover, with $H_\alpha \in \mathfrak{h}$ as in §2.3,

$$\mathfrak{h}_{\mathbb{Z}[a]} \;:=\; Span_{\mathbb{Z}[a]}\big(H_1, H_2, H_3\big) \;=\; Span_{\mathbb{Z}[a]}\big(\{H_\alpha \,|\, \alpha\in\Delta\}\big) \quad ;$$

(b) $\big[H_i\,, H_j\big] = 0\,, \quad \big[H_i\,, X_\alpha\big] = \alpha(H_i)\,X_\alpha\,, \qquad \forall\; i,j \in \{1,\dots,\ell\}\,,\; \alpha\in\Delta\;;$

(c) $\big[X_\alpha\,, X_{-\alpha}\big] = \sigma_\alpha\, H_\alpha \qquad \forall\;\alpha\in\Delta$

with H_α as in *(a)*, and $\sigma_\alpha := -1$ if $\alpha \in \Delta_1^-$, $\sigma_\alpha := 1$ otherwise;

(d) $\big[X_\alpha\,,\,X_\beta\big] \,=\, c_{\alpha,\beta}\,X_{\alpha+\beta} \quad \forall\;\alpha,\beta\in\Delta\,:\,\alpha+\beta\neq 0\,,$ with

(d.1) if $(\alpha+\beta)\notin\Delta\,,$ then $c_{\alpha,\beta}=0\,,$ and $X_{\alpha+\beta}:=0\,,$

(d.2) if $\big(\alpha,\alpha\big)\neq 0$ or $\big(\beta,\beta\big)\neq 0\,,$ and if $\Sigma^\alpha_\beta := \big(\beta+\mathbb{Z}\,\alpha\big)\cap\big(\Delta\cup\{0\}\big) = \big\{\beta-r\,\alpha\,,\,\dots\,,\,\beta+q\,\alpha\big\}$ is *the α–string through β*, then $c_{\alpha,\beta}=\pm(r+1)\,;$

(d.3) if $\big(\alpha\,,\alpha\big) \,=\, 0 = \big(\beta\,,\beta\big),$ then $c_{\alpha,\beta}=\pm\beta\big(H_\alpha\big)\,.$

Definition 3.2. If B is a Chevalley basis of $\mathfrak{g}$, we denote by $\mathfrak{g}^{\mathbb{Z}[a]}$ the $\mathbb{Z}[a]$–span of B, and we call it the *Chevalley Lie superalgebra* (of $\mathfrak{g}$).

3.2 Existence of Chevalley bases

The *existence* of Chevalley bases (with slightly different definition) is proved in [7]; we now present an explicit example of such a basis.

Let us consider in $\mathfrak{g}$ the generators $h_i\,,\ e_i\,,\ f_i\ \ (\,i=1,2,3\,)$ and the root vectors $e_{1,2}\,,\ e_{1,3}\,,\ e_{1,2,3}\,,\ e_{1,1,2,3}\,,\ f_{2,1}\,,\ f_{3,1}\,,\ f_{3,2,1}\,,\ f_{3,2,1,1}$ constructed in §2.3. Looking at all brackets among them considered there, it is a routine matter to check that the set

$$B \,:=\, \big\{H_i\,,\,e_i\,,\,f_i\big\}_{i=1,2,3}\bigcup\big\{e_{1,2}\,,e_{1,3}\,,e_{1,2,3}\,,e_{1,1,2,3}\,,f_{2,1}\,,f_{3,1}\,,f_{3,2,1}\,,f_{3,2,1,1}\big\}$$

with $H_1:=h_1\,,\ \ H_2:=(1+a)^{-1}\big(2h_1-h_2-ah_3\big)\,,\ \ H_3:=h_3\,,$ is indeed a *Chevalley basis for $\mathfrak{g}=D(2,1;a)$* — the proof is a bookkeeping matter.

To be precise, the root vectors in B are:

— *even*: $X_{+(2\,\alpha_1+\alpha_2+\alpha_3)} = e_{1,1,2,3}\,,\ \ X_{+\alpha_2}=e_2\,,\ \ X_{+\alpha_3}=e_3$ *(positive)*
$X_{-(2\,\alpha_1+\alpha_2+\alpha_3)} = f_{3,2,1,1}\,,\ \ X_{-\alpha_2}=f_2\,,\ \ X_{-\alpha_3}=f_3$ *(negative)*
— *odd*: *(positive)* $X_{+\alpha_1}=e_1\,,\ \ X_{+\alpha_1+\alpha_2}=e_{1,2}\,,\ \ X_{+\alpha_1+\alpha_3}=e_{1,3}\,,$
$X_{+\alpha_1+\alpha_2+\alpha_3}=e_{1,2,3}$
(negative) $X_{-\alpha_1}=f_1\,,\ \ X_{-\alpha_1-\alpha_2}=f_{2,1}\,,\ \ X_{-\alpha_1-\alpha_3}=e_{3,1}\,,$
$X_{-\alpha_1-\alpha_2-\alpha_3}=f_{3,2,1}$

while the Cartan elements in B are just $H_1\,,\ H_2$ and H_3 defined as above.

3.3 Kostant superalgebra

For any $\mathbb{K}$–algebra A, given $n \in \mathbb{N}$ and $y \in A$ we define the *n–th binomial coefficients* $\binom{y}{n}$ and the *n–th divided power* $y^{(n)}$ by

$$\binom{y}{n} := \frac{y(y-1)\cdots(y-n+1)}{n!} \quad , \qquad y^{(n)} := y^n\big/n!$$

Recall that $\mathbb{Z}[a]$ is the unital subring of $\mathbb{K}$ generated by a. We need also to consider the ring $\mathbb{Z}_a$, which by definition is the unital subring of $\mathbb{K}$ generated by the subset $\left\{ \binom{P(a)}{n} \,\middle|\, P(a) \in \mathbb{Z}[a]\,,\, n \in \mathbb{N} \right\}$.

By a classical result on integer-valued polynomials (see [4], Lemma 4.1) one shows that $\mathbb{Z}_a$ in fact is generated by $\left\{ \binom{a}{n} \,\middle|\, n \in \mathbb{N} \right\}$ as well. Note also that $\mathbb{Z}_a = \mathbb{Z}$ if and only if $a \in \mathbb{Z}$.

Fix in $\mathfrak{g} := D(2,1;a)$ a Chevalley basis $B = \{H_1, H_2, H_3\} \coprod \{X_\alpha\}_{\alpha\in\Delta}$ as in Definition 3.1. Let $U(\mathfrak{g})$ be the universal enveloping superalgebra of $\mathfrak{g}$.

In [4], §4.1, the Kostant superalgebra $K_{\mathbb{Z}}(\mathfrak{g})$ was defined as the subalgebra of $U(\mathfrak{g})$ generated by special elements: divided powers of the root vectors attached to even roots, root vectors attached to odd roots, and binomial coefficients in the elements of $\mathfrak{h}_{\mathbb{Z}}$, the $\mathbb{Z}$–span of the elements of $\{H_1, H_2, H_3\}$.

If we try to perform the same construction *verbatim* for $\mathfrak{g} = D(2,1;a)$ for $a \in \mathbb{K} \setminus \mathbb{Z}$, we are soon forced to include among the generators all binomial coefficients of type $\binom{H}{n}$ with $H \in \mathfrak{h}_{\mathbb{Z}[a]}$, the $\mathbb{Z}[a]$–span of $\{H_1, H_2, H_3\}$. When commuting such binomial coefficients with divided powers, coefficients of type $\binom{\alpha(H)}{n}$ show up, where α is a root and $H \in \mathfrak{h}_{\mathbb{Z}[a]}$. By construction we have $\alpha(H) \in \mathbb{Z}[a]$, hence $\binom{\alpha(H)}{n}$ belongs to the ring $\mathbb{Z}_a$ defined above.

In force of the above remarks, for $\mathfrak{g} := D(2,1;a)$ we define the Kostant superalgebra $K_{\mathbb{Z}_a}(\mathfrak{g})$ like we did in [4] for the other classical Lie superalgebras, *but* with $\mathbb{Z}_a$ replacing $\mathbb{Z}$ as ground ring: more precisely, we set

Definition 3.3. We call *Kostant superalgebra*, or Kostant's $\mathbb{Z}_a$–form of $U(\mathfrak{g})$, the unital $\mathbb{Z}_a$–subsuperalgebra $K_{\mathbb{Z}_a}(\mathfrak{g})$ of $U(\mathfrak{g})$ generated by

$$X_\alpha^{(n)} := X_\alpha^n\big/n!\ , \quad X_\gamma\ , \quad \binom{H}{n} \qquad \forall\ \alpha \in \Delta_0\ ,\ n \in \mathbb{N}\,,\ \gamma \in \Delta_1\,,\ H \in \mathfrak{h}_{\mathbb{Z}[a]}$$

The following analogue of Corollary 4.2 in [4] holds (with same proof), which is needed in the proof of the PBW-like theorem for $K_{\mathbb{Z}_a}(\mathfrak{g})$:

Corollary 3.4.

(a) $\mathbb{H}_{\mathbb{Z}_a} := \big\{\, h \in U(\mathfrak{h}) \;\big|\; h\big(z'_1, z'_2, z'_3\big) \in \mathbb{Z}_a ,\ \ \forall\, z'_1, z'_2, z'_3 \in \mathbb{Z}_a \,\big\}$ *is a free* $\mathbb{Z}_a$*–submodule of* $U(\mathfrak{h})$*, with basis* $B_{U(\mathfrak{h})} := \Big\{ \binom{H_1}{n_1} \binom{H_2}{n_2} \binom{H_3}{n_3} \,\Big|\, n_1, n_2, n_3 \in \mathbb{N} \Big\}$.

(b) The $\mathbb{Z}_a$*–subalgebra of* $U(\mathfrak{g})$ *generated by all the elements* $\binom{H'-z'}{n}$ *with* $H' \in \mathfrak{h}_{\mathbb{Z}_a} := \sum_{i=1}^{3} \mathbb{Z}_a H_i$, $z' \in \mathbb{Z}_a$, $n \in \mathbb{N}$ *, is nothing but* $\mathbb{H}_{\mathbb{Z}_a}$.

3.4 Commutation rules and Kostant's theorem

In [4] the authors proved a "super PBW-like" theorem for the Kostant's superalgebra: namely, the latter is a free $\mathbb{Z}$–module with $\mathbb{Z}$–basis the set of ordered monomials (w. r. to any total order) whose factors are binomial coefficients in the H_i's, or *odd* root vectors, or divided powers of *even* root vectors. This result follows from a direct analysis of commutation rules among the generators of the Kostant's superalgebra. The same procedure will work for $D(2,1;a)$, once such commutation rules are established.

We list hereafter all such rules (and a few more general ones too); their most relevant feature is that all coefficients in these relations are in $\mathbb{Z}_a$. All of them are proved via simple induction arguments (left to the reader).

We split the list into two sections: *(1)* relations involving only *even* generators (known by classical theory); *(2)* relations involving also *odd* generators.

(1) Even generators only (that is $\binom{H_i}{m}$'s and $X_\alpha^{(n)}$'s only, $\alpha \in \Delta_0$)**:**

$$\begin{aligned}
\binom{H_i}{n}\binom{H_j}{m} &= \binom{H_j}{m}\binom{H_i}{n} && \forall\ i,j \in \{1,2,3\}\,,\quad \forall\ n,m \in \mathbb{N} \\
X_\alpha^{(n)}\, f(H) &= f\big(H - n\,\alpha(H)\big)\, X_\alpha^{(n)} && \forall\ \alpha \in \Delta_0\,,\ H \in \mathfrak{h}\,, \\
& && \forall\ n \in \mathbb{N}\,,\ f(T) \in \mathbb{K}[T] \\
X_\alpha^{(n)}\, X_\alpha^{(m)} &= \binom{n+m}{m} X_\alpha^{(n+m)} && \forall\, \alpha \in \Delta_0\,,\ \forall\ n,m \in \mathbb{N} \\
X_\alpha^{(n)}\, X_\beta^{(m)} &= X_\beta^{(m)}\, X_\alpha^{(n)} + \ l.h.t && \forall\, \alpha,\beta \in \Delta_0\,,\ \forall\ n,m \in \mathbb{N}
\end{aligned}$$

where *l.h.t.* stands for a $\mathbb{Z}_a$–linear combinations of monomials in the $X_\delta^{(k)}$'s and in the $\binom{H_i}{c}$'s whose "height" — by definition, the sum of all "exponents" k occurring in such a monomial — is less than $n+m$. A *special case* is

$$X_\alpha^{(n)}\, X_{-\alpha}^{(m)} \;=\; \sum_{k=0}^{min(m,n)} X_{-\alpha}^{(m-k)} \binom{H_\alpha - m - n + 2\,k}{k} X_\alpha^{(n-k)} \qquad \forall\; \alpha \in \Delta_0\,,\;\; \forall\;\, m, n \in \mathbb{N}$$

(2) Odd and even generators (also involving the X_γ's, $\gamma \in \Delta_1$)**:**

$$\begin{aligned}
X_\gamma\, f(H) \;&=\; f\big(H - \gamma(H)\big)\, X_\gamma && \forall\;\, \gamma \in \Delta_1\,,\; h \in \mathfrak{h}\,,\; f(T) \in \mathbb{K}[T] \\
X_\gamma^n \;&=\; 0 && \forall\;\, \gamma \in \Delta_1\,, \qquad \forall\;\, n \geq 2 \\
X_{-\gamma}\, X_\gamma \;&=\; -X_\gamma\, X_{-\gamma} \;+\; H_\gamma && \forall\;\, \gamma \in \Delta_1 \\
X_\gamma\, X_\delta \;&=\; -X_\delta\, X_\gamma \;+\; c_{\gamma,\delta}\, X_{\gamma+\delta} && \forall\;\, \gamma, \delta \in \Delta_1\,,\; \gamma + \delta \neq 0
\end{aligned}$$

with $c_{\gamma,\delta}$ as in Definition 3.1,

$$X_\alpha^{(n)}\, X_\gamma \;=\; X_\gamma\, X_\alpha^{(n)} \;+\; \sum_{k=1}^{n} \Big(\prod_{s=1}^{k} \varepsilon_s\Big) \binom{r+k}{k} X_{\gamma+k\,\alpha}\, X_\alpha^{(n-k)} \qquad \forall\;\, n \in \mathbb{N}\,, \quad \forall\;\, \alpha \in \Delta_0\,,\; \gamma \in \Delta_1$$

with $\sigma_\gamma^\alpha = \big\{\gamma - r\,\alpha\,, \dots, \gamma\,, \dots, \gamma + q\,\alpha\big\}$, $X_{\gamma+k\,\alpha} := 0$ if $(\gamma + k\,\alpha) \notin \Delta$,
and $\varepsilon_s = \pm 1$ such that $\big[X_\alpha\,, X_{\gamma+(s-1)\,\alpha}\big] = \varepsilon_s\,(r+s)\, X_{\gamma+s\,\alpha}$

Here now is our super-version of Kostant's theorem for $K_{\mathbb{Z}_a}(\mathfrak{g})$:

Theorem 3.5. *The Kostant superalgebra $K_{\mathbb{Z}_a}(\mathfrak{g})$ is a free $\mathbb{Z}_a$–module. For any given total order $\preceq$ of the set $\Delta \cup \{1\,,2\,,3\}$, a $\mathbb{Z}_a$–basis of $K_{\mathbb{Z}_a}(\mathfrak{g})$ is the set $\mathcal{B}$ of ordered "PBW-like monomials", i.e. all products (without repetitions) of factors of type $X_\alpha^{(n_\alpha)}$, $\binom{H_i}{n_i}$ and X_γ — with $\alpha \in \Delta_0$, $i \in \{1\,,2\,,3\}$, $\gamma \in \Delta_1$ and n_α, $n_i \in \mathbb{N}$ — taken in the right order with respect to $\preceq$.*

This result is proved exactly like the similar one in [4], making use of the commutation relations considered above. It also has a direct consequence, again proved like in [4]. To state it, let first consider $\mathfrak{g}_1^{\mathbb{Z}[a]}$, the odd part of $\mathfrak{g}^{\mathbb{Z}[a]}$: it has $\big\{X_\gamma \,\big|\, \gamma \in \Delta_1\big\}$ as $\mathbb{Z}[a]$–basis, by construction. Then let $\bigwedge \mathfrak{g}_1^{\mathbb{Z}[a]}$ be the exterior $\mathbb{Z}[a]$–algebra over $\mathfrak{g}_1^{\mathbb{Z}[a]}$, and let $\bigwedge \mathfrak{g}_1^{\mathbb{Z}_a}$ be its scalar extension to $\mathbb{Z}_a$. Let also $K_{\mathbb{Z}}(\mathfrak{g}_0)$ be the classical Kostant's algebra of $\mathfrak{g}_0$ (over $\mathbb{Z}$) and let $K_{\mathbb{Z}_a}(\mathfrak{g}_0)$ be its scalar extension to $\mathbb{Z}_a$. Then the tensor factorization $U(\mathfrak{g}) \cong U(\mathfrak{g}_0) \otimes_{\mathbb{K}} \bigwedge \mathfrak{g}_1$ (see [14]) has the following "integral version":

Corollary 3.6. *There exists an isomorphism of $\mathbb{Z}_a$–modules*

$$K_{\mathbb{Z}_a}(\mathfrak{g}) \;\cong\; K_{\mathbb{Z}_a}(\mathfrak{g}_0) \otimes_{\mathbb{Z}_a} \bigwedge \mathfrak{g}_1^{\mathbb{Z}_a}$$

Remarks 3.7.

(a) Following a classical pattern (and cf. [1], [2], [12] in the super context) we can define the *superalgebra of distributions* $\mathcal{D}ist(G)$ on any supergroup G . Then $\mathcal{D}ist(G) = K_{\mathbb{Z}_a}(\mathfrak{g}) \otimes_{\mathbb{Z}_a} \mathbb{K}$, when $\mathfrak{g} := Lie(G)$ is just $D(2,1;a)$.

(b) In this section we proved that the assumptions of Theorem 2.8 in [12] do hold for any supergroup G whose tangent Lie superalgebra is just $\mathfrak{g} = D(2,1;a)$. Therefore, *all results in [12] do apply to such supergroups.*

4 Chevalley supergroups of type $D(2,1;a)$

Classically, Chevalley groups are defined as follows. Let $\mathfrak{g}$ be a finite dimensional semisimple Lie algebra over an algebraically closed field $\mathbb{K}$ of characteristic zero. Choosing a Chevalley basis of $\mathfrak{g}$, we can define a Kostant form $K_{\mathbb{Z}}(\mathfrak{g})$, generated by divided powers of root vectors. Then any simple $\mathfrak{g}$–module V contains a $\mathbb{Z}$–lattice M, which is $K_{\mathbb{Z}}(\mathfrak{g})$–stable, hence $K_{\mathbb{Z}}(\mathfrak{g})$ acts on M . Using this action and its scalar extension to a field $\Bbbk$, we consider, inside the group $\mathrm{GL}(V_{\Bbbk})$, $V_{\Bbbk} := \Bbbk \otimes_{\mathbb{Z}} M$, additive one-parameter subgroups associated to all root vectors and multiplicative one-parameter subgroups associated to all coroots. The Chevalley group (associated to $\mathfrak{g}$ and V) is then the subgroup of $\mathrm{GL}(V_{\Bbbk})$ generated by such one-parameter subgroups.

Now we provide a similar construction for the Lie superalgebra $\mathfrak{g} = D(2,1;a)$, suitably adapting what is done in [4] for the other classical Lie superalgebra (including $\mathfrak{g} = D(2,1;a)$ when the parameter a is an integer).

4.1 Admissible lattices

Let $\mathbb{K}$ be an algebraically closed field of characteristic zero. If R is a unital subring of $\mathbb{K}$, and V a finite dimensional $\mathbb{K}$–vector space, any $M \subseteq V$ is called *R–lattice* (or R–form) of V if $M = Span_R(B)$ for some basis B of V .

Let $\mathfrak{g} = D(2,1;a)$ be defined over $\mathbb{K}$, and fix the ring $\mathbb{Z}_a$, a Chevalley basis B of $\mathfrak{g}$ and the Kostant algebra $K_{\mathbb{Z}_a}(\mathfrak{g})$ be as in §3.

Definition 4.1. Let V be a $\mathfrak{g}$–module, and let M be a $\mathbb{Z}_a$–lattice of it.

(a) We call V *rational* if:

(a-1) $\mathfrak{h}_{\mathbb{Z}[a]} := \mathit{Span}_{\mathbb{Z}[a]}\big(H_1, H_2, H_3\big)$ acts diagonally on V with eigenvalues in $\mathbb{Z}[a]$; in other words, one has $V = \bigoplus_{\mu\in\mathfrak{h}^*} V_\mu$, with $V_\mu := \big\{ v \in V \,\big|\, h.v = \mu(h)\, v \;\forall\, h \in \mathfrak{h} \big\}$, *and* $\mu(H_i) \in \mathbb{Z}[a]$ for all i and all μ such that $V_\mu \neq \{0\}$;

(a-2) $\mu(H_\alpha) \in \mathbb{Z}$ for all $\alpha \in \Delta_0$ and $\mu \in \mathfrak{h}^*$ such that $V_\mu \neq \{0\}$.

(b) We call M *admissible (lattice)* if it is $K_{\mathbb{Z}_a}(\mathfrak{g})$–stable.

The above is a very slight variation of the analogue definition in [4], §5.1. Similarly, the next two results are proved like there:

Theorem 4.2. *Any rational, finite dimensional, semisimple $\mathfrak{g}$–module V contains an admissible lattice M. Any such M is the (direct) sum of its weight components, i.e.* $M = \bigoplus\limits_{\mu\in\mathfrak{h}^*} M_\mu$ *with* $M_\mu = M \cap V_\mu$.

Theorem 4.3. *Let V be a rational, finite dimensional $\mathfrak{g}$–module, M an admissible lattice of V, and* $\mathfrak{g}_V = \big\{ X \in \mathfrak{g} \,\big|\, X.M \subseteq M \big\}$. *If V is faithful, then*

$$\mathfrak{g}_V \;=\; \mathfrak{h}_V \bigoplus \big(\oplus_{\alpha\in\Delta} \mathbb{Z}_a X_\alpha \big) \;, \qquad \mathfrak{h}_V \;:=\; \big\{ H \in \mathfrak{h} \;\big|\; \mu(H) \in \mathbb{Z}_a \,,\; \forall\, \mu \in \Lambda \big\}$$

where Λ is the set of all weights of V. In particular, $\mathfrak{g}_V$ is a $\mathbb{Z}_a$–lattice in $\mathfrak{g}$, independent of the choice of the admissible lattice M (but not of V).

From now on, we retain the following notation: V is a rational, finite dimensional $\mathfrak{g}$–module, and M is an admissible lattice of V.

Like in §2.2, we assume $\Bbbk$ to be a commutative unital ring such that 2 and 3 are not zero and they are not zero divisors in $\Bbbk$. In addition, now

we assume $\Bbbk$ to be also a (commutative unital) $\mathbb{Z}_a$–algebra $\Bbbk$.

With these assumptions, we set

$$\mathfrak{g}_\Bbbk \;:=\; \Bbbk \otimes_{\mathbb{Z}_a} \mathfrak{g}_V \;\;, \qquad V_\Bbbk \;:=\; \Bbbk \otimes_{\mathbb{Z}_a} M \;\;, \qquad U_\Bbbk(\mathfrak{g}) \;:=\; \Bbbk \otimes_{\mathbb{Z}_a} K_{\mathbb{Z}_a}(\mathfrak{g}) \;\;;$$

then $\mathfrak{g}_\Bbbk$ acts faithfully on $V_\Bbbk$, which yields an embedding of $\mathfrak{g}_\Bbbk$ into $\mathfrak{gl}(V_\Bbbk)$.

For any $A \in (\mathrm{salg})_\Bbbk = (\mathrm{salg})$, the Lie superalgebra $\mathfrak{g}_A := A \otimes_\Bbbk \mathfrak{g}_\Bbbk$ acts faithfully on $V_\Bbbk(A) := A \otimes_\Bbbk V_\Bbbk$, hence it embeds into $\mathfrak{gl}\big(V_\Bbbk(A)\big)$, etc.

4.2 Additive one-parameter supersubgroups

Let $\alpha \in \Delta_0$, $\beta \in \Delta_1$, and let X_α , X_β be the associated root vectors (in our fixed Chevalley basis of $\mathfrak{g}$). Both X_α and X_β act as nilpotent operators on V , hence on M and $V_{\Bbbk}$, so they are represented by nilpotent matrices in $\mathfrak{gl}(V_{\Bbbk}(A))$; the same holds for all operators

$$t\,X_\alpha\,,\ \ \vartheta\,X_\beta\ \in\ \mathrm{End}\big(V_{\Bbbk}(A)\big) \qquad\qquad \forall\ t \in A_0\,,\ \vartheta \in A_1\ . \tag{4.1}$$

We see at once that, for any $n \in \mathbb{N}$, we have $Y^{(n)} := Y^n\big/n! \in \big(K_{\mathbb{Z}}(\mathfrak{g})\big)(A)$ for any Y as in (4.1); moreover, $Y^{(n)} = 0$ for $n \gg 0$, by nilpotency. Thus the formal power series $\exp(Y) := \sum_{n=0}^{+\infty} Y^{(n)}$, when computed for Y as in (4.1), gives a well-defined element in $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$, expressed as finite sum.

Definition 4.4. Let $\alpha \in \Delta_0$, $\beta \in \Delta_1$, and X_α , X_β as above. We define the supergroup functors x_α and x_β from (salg) to (groups) as

$$x_\alpha(A) := \big\{ x_\alpha(t) := \exp\big(tX_\alpha\big) \,\big|\, t \in A_0 \big\} \ =\ \big\{ \big(1 + t\,X_\alpha + \tfrac{t}{2!}\,X_\alpha^{\,2} + \cdots\big) \,\big|\, t \in A_0 \big\}$$

$$x_\beta(A) := \big\{ x_\beta(\vartheta) := \exp\big(\vartheta X_\beta\big) \,\big|\, \vartheta \in A_1 \big\} \ =\ \big\{ \big(1 + \vartheta\,X_\beta\big) \,\big|\, \vartheta \in A_1 \big\}$$

For later convenience we shall also formally write $x_\zeta(\mathbf{t}) := 1$ when ζ belongs to the $\mathbb{Z}$–span of Δ but $\zeta \not\in \Delta$.

Like in [4], Proposition 4.5*(a)*, one sees that these supergroup functors are in fact representable, hence they both are *affine supergroups*: namely, x_α is represented by $\Bbbk[x]$ and x_β by $\Bbbk[\xi]$, these being Hopf algebras with coproduct $\Delta(x) = x \otimes 1 + 1 \otimes x$ and $\Delta(\xi) = \xi \otimes 1 + 1 \otimes \xi$. We shall refer to both x_α and x_β as *additive one-parameter (super)subgroups*.

4.3 Multiplicative one-parameter supersubgroups of classical type

Consider $\mathfrak{h}_{\mathbb{Z}}^0 := \mathit{Span}_{\mathbb{Z}}\big(\big\{H_\alpha \,\big|\, \alpha \in \Delta_0\big\}\big)$; this is a $\mathbb{Z}$–form of $\mathfrak{h}$, with $\mathfrak{h}_{\mathbb{Z}}^0 \subseteq \mathfrak{h}_{\mathbb{Z}[a]} := \mathit{Span}_{\mathbb{Z}[a]}\big(H_1, H_2, H_3\big)$. For any $\alpha \in \Delta_0$ $\big(\subseteq \mathfrak{h}^*\big)$, let $H_\alpha \in \mathfrak{h}_{\mathbb{Z}}^0$ be the corresponding coroot (cf. §2.3). Let $V = \oplus_\mu V_\mu$ be the splitting of V into weight spaces; as V is rational, we have $\mu(H_\alpha) \in \mathbb{Z}$ for all $\alpha \in \Delta_0$ and $\mu \in \mathfrak{h}^* : V_\mu \neq \{0\}$. Now, for any $A \in$ (salg) , $\alpha \in \Delta_0$ and $t \in U(A_0)$ (the group of invertible elements in A_0) set $h_\alpha(t).v := t^{\mu(H_\alpha)}\, v$ for all $v \in V_\mu$, $\mu \in \mathfrak{h}^*$: this defines another operator (also locally expressed by exponentials)

$$h_\alpha(t) \in \mathrm{GL}\big(V_{\Bbbk}(A)\big) \qquad\qquad \forall\ t \in U(A_0)\,,\ \alpha \in \Delta_0\ . \tag{4.2}$$

More in general, if $\{H_{\alpha_i}\}_{i=1,2,3}$ is any basis of Δ_0 and $H = \sum_{i=1}^{3} z_i H_{\alpha_i}$ (with $z_1, z_2, z_3 \in \mathbb{Z}$) then we define $h_H(t) := \prod_{i=1}^{3} \big(h_{\alpha_i}(t)\big)^{z_i}$, for $\alpha \in \Delta_0$.

Definition 4.5. Let $H \in \mathfrak{h}_{\mathbb{Z}}^0$ as above. We define the supergroup functor h_H — also writing $h_\alpha := h_{H_\alpha}$ for any $\alpha \in \Delta_0$ — from (salg) to (groups) as

$$A \ \mapsto\ h_H(A) \ :=\ \big\{\, t^H := h_H(t) \;\big|\; t \in U(A_0) \big\}$$

As in [4], Proposition 4.5*(b)*, one sees that these functors are representable, so they are affine supergroups; even more, they are also closed subgroups of the diagonal subgroup of $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$.

4.4 Multiplicative one-parameter supersubgroups of a–type

In order to attach a suitable "multiplicative one-parameter supersubgroups" to *any* Cartan element in $\mathfrak{h}_{\mathbb{Z}[a]} := \mathit{Span}_{\mathbb{Z}[a]}\big(H_1, H_2, H_3\big)$ — not only in $\mathfrak{h}_{\mathbb{Z}}^0$ as above — we need to suitably adapt our previous construction.

Consider a Cartan element H_i in our fixed Chevalley basis: we want to define a suitable, representable supergroup functor associated to it, to be called $h_i^{[a]}$. Given any $A \in$ (salg) , and $t \in U(A_0)$, we look for an operator like $h_i^{[a]}(t) := t^{H_i} \in \mathrm{GL}\big(V_{\Bbbk}(A)\big)$. By formal identities, it should be given by

$$h_i^{[a]}(t) \ :=\ t^{H_i} \ =\ \big(1 + (t-1)\big)^{H_i} \ =\ \textstyle\sum_{n=0}^{+\infty} (t-1)^n \binom{H_i}{n}$$

Let $V = \bigoplus_{\mu \in \mathfrak{h}^*} V_\mu$ be the splitting of V into weight spaces; definitions imply

$$h_i^{[a]}(t)\Big|_{V_\mu} \ =\ \textstyle\sum_{n=0}^{+\infty} (t-1)^n \binom{\mu(H_i)}{n}\, id_{V_\mu} \ =\ t^{\mu(H_i)}\, id_{V_\mu}$$

on weight spaces, which makes sense — and then globally yields a well-defined operator on all of $V = \bigoplus_{\mu \in \mathfrak{h}^*} V_\mu$ — as soon as $t^{\mu(H_i)} := \sum_{n=0}^{+\infty} (t-1)^n \binom{\mu(H_i)}{n}$

is a well-defined element of A. Now, as V is rational and M is admissible, we have $\mu(H_i) \in \mathbb{Z}[a]$. Therefore, a necessary condition we may require is

$$t^{z(a)} := \textstyle\sum_{n=0}^{+\infty} (t-1)^n \binom{z(a)}{n} \in A \qquad \forall\ z(a) \in \mathbb{Z}[a] \tag{4.3}$$

which in the end is equivalent to

$$t^{\pm a^k} := \textstyle\sum_{n=0}^{+\infty} (t-1)^n \binom{\pm a^k}{n} \in A \qquad \forall\ k \in \mathbb{N} \tag{4.4}$$

Both (4.3) and (4.4) must be read as conditions defining a suitable subset of A_0, namely that of all elements $t \in A_0$ for which the condition does hold.

Now we go and fix details, as to give a well-defined meaning to the formal series expressions in (4.3–4) and to the just sketched construction.

Consider the polynomial $\mathbb{Z}_a$–superalgebra $\mathbb{Z}_a[\ell]$, where ℓ is an *even* indeterminate: this is also a (super)bialgebra, with $\Delta(\ell) = \ell \otimes \ell$, $\epsilon(\ell) = 1$. Let $\mathbb{Z}_a[[\ell-1]]$ be the $(\ell-1)$–adic completion of $\mathbb{Z}_a[\ell]$; this contains also an inverse of ℓ, namely $\ell^{-1} = \sum_{n=0}^{+\infty} (-1)^n (\ell-1)^n$, so $\mathbb{Z}_a\big[\ell, \ell^{-1}\big] \subseteq \mathbb{Z}_a[[\ell-1]]$. The coproduct of $\mathbb{Z}_a[\ell]$ extends to $\mathbb{Z}_a[[\ell-1]]$ making it into a *formal* bialgebra — the coproduct taking values into the $(\ell-1)$–adic completion of the algebraic tensor square of $\mathbb{Z}_a[[\ell-1]]$ — which indeed is a *formal Hopf algebra*. In the latter, both ℓ and ℓ^{-1} are group-like elements, so that $\mathbb{Z}_a\big[\ell, \ell^{-1}\big]$ is a (non formal) Hopf subalgebra of $\mathbb{Z}_a[[\ell-1]]$. Consider the elements

$$\ell^{\pm a^k} := \textstyle\sum_{n=0}^{+\infty} (\ell-1)^n \binom{\pm a^k}{n} \in \mathbb{Z}_a[[\ell-1]] \qquad\qquad \forall\ k \in \mathbb{N}$$

A key property of these elements is the following:

Lemma 4.6. *All the $\ell^{\pm a^k}$'s ($k \in \mathbb{N}$) are group-like elements of $\mathbb{Z}_a[[\ell-1]]$.*

Proof. When ℓ ranges in the complex field $\mathbb{C}$ and $a \in \mathbb{C} \setminus \{0, -1\}$, the power series $\sum\limits_{n=0}^{+\infty} (\ell-1)^n \binom{\pm a^k}{n}$ represents the Taylor expansion of the analytic function $\phi_a : \ell \mapsto \ell^{\pm a^k}$ in a neighbourhood of $\ell_0 := 1$. Now, the function ϕ_a is multiplicative, i.e. $\phi_a\big(\ell_1 \ell_2\big) = \phi_a(\ell_1)\, \phi_a(\ell_2)$; this identity for all complex values of ℓ_1 and ℓ_2 in a neighbourhood of 1 implies (passing through Taylor expansion) a similar identity at the level of power series. In turn, the latter identity implies an identity among *formal power series* (i.e., still holding when complex numbers ℓ_1, ℓ_2 are replaced with indeterminates). This can be recast saying that the formal power series $\sum_{n=0}^{+\infty} (\,\ell-1)^n \binom{\pm a^k}{n}$ is a group-like element in $\mathbb{Z}_a[[\,\ell-1]]$. As this holds for *any* $a \in \mathbb{C}$, it must hold for a formal parameter a, i.e. if the complex value of a is replaced by an indeterminate. Then the formal symbol a can be replaced by any genuine value in $\mathbb{K} \setminus \{0,-1\}$, and the formal series in $\mathbb{Z}_a[[\,\ell-1]]$ will still be group-like. $\square$

Let now $\mathbb{Z}_a\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$ be the $\mathbb{Z}_a$–subalgebra of $\mathbb{Z}_a[[\,\ell-1]]$ generated by all the $\ell^{\pm a^k}$'s: as these generators are group-like, $\mathbb{Z}_a\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$ is in fact a Hopf sub(super)algebra of $\mathbb{Z}_a[[\ell-1]]$. In particular, it is a (totally even) Hopf algebra over $\mathbb{Z}_a$. If now $\Bbbk$ is any (unital, commutative) $\mathbb{Z}_a$–algebra as above, then $\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big] := \Bbbk \otimes_{\mathbb{Z}_a} \mathbb{Z}_a\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$ belongs to $(\mathrm{salg})_\Bbbk = (\mathrm{salg})$, and in addition it is a Hopf algebra over $\Bbbk$.

Given $A \in (\mathrm{salg})$, any algebra morphism φ from $\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$ to A is uniquely determined by the choice of a single element $t \in A$, namely $t = \varphi(\ell)$; in particular, $t \in A_0$ — as ℓ is even — and $t \in U(A_0)$ — since ℓ is invertible. Therefore, we can define

$$P_a(A) := \Big\{\, t \in U(A_0) \;\Big|\; \exists\ \varphi \in \mathrm{Hom}\Big(\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big], A\Big) \;:\; \varphi(\ell) = t \,\Big\} \qquad (4.5)$$

By construction, this is a group. Indeed, this construction defines a representable functor $P_a : (\mathrm{salg}) \longrightarrow (\mathrm{groups})$, given on objects by $A \mapsto P_a(A)$ and represented by $\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$. As the latter is a Hopf $\Bbbk$–(super)algebra, the functor P_a is indeed an affine supergroup functor. Actually, P_a is just a generalization of the functor $A \mapsto U_0(A) := U(A_0)$, which is represented by $\Bbbk\big[\ell, \ell^{-1}\big]$. Indeed, one has $P_a = U_0$ if and only if $a \in \mathbb{Z}$.

Remarks 4.7.

(a) By (4.4) it is easy to see — using notation of §2 — that for any $A \in (\mathrm{salg})$ one has $P_a(A) \supseteq \big(1 + \mathfrak{N}(A_0)\big) \supseteq \big(1 + A_1^2\big)$.

(b) It is clear that $P_a(\mathbb{C}) = \mathbb{C}^* = U(\mathbb{C})$, with $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. Now assume $\mathbb{K} = \mathbb{C}$ and $A = \mathbb{C}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]$, where the x_i's and the ξ_j's respectively are even and odd indeterminates. If $(\xi_1, \dots, \xi_n)$ is the ideal generated by the ξ_i's, one has $P_a\big(\mathbb{C}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]\big) = \mathbb{C}^* + (\xi_1, \dots, \xi_n)^2$. In particular, in this case one even has

$$P_a\big(\mathbb{C}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]\big) \;=\; U\big(\mathbb{C}[x_1, \dots, x_m, \xi_1, \dots, \xi_n]_0\big)$$

By the same argument, one has also $P_a(A) = U(A_0)$ for all $A \in (\mathrm{salg})_{\mathbb{C}}$ such that $U(A_0) = \mathbb{C}^* + \mathfrak{N}(A_0)$, where $\mathfrak{N}(A_0)$ is the nilradical of A_0. $\diamondsuit$

Now let $V = \bigoplus_{\mu\in\mathfrak{h}^*} V_\mu$ and M be as above. For any $A \in (\text{salg})$ and any $t \in P_a(A)$, the expressions in (4.3–4) do make sense, i.e. they provide unique, well-defined elements of A, namely $t^{z(a)} := \varphi\big(\ell^{z(a)}\big)$ — for (4.3), say — with φ like in (4.5). Therefore, the formula

$$h_i^{[a]}(t) \,:=\, t^{H_i} \,=\, \sum_{n=0}^{+\infty} (t-1)^n \binom{H_i}{n} \,\in\, \mathrm{GL}\big(V_{\Bbbk}(A)\big) \qquad \forall\ t \in P_a(A)$$

(for $i = 1,2,3$) yields an element of $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$ whose action is described by $h_i^{[a]}(t).v \,:=\, t^{\mu(H_i)}\, v$ for all $v \in V_\mu$, $\mu \in \mathfrak{h}^* : V_\mu \neq \{0\}$ ($i = 1,2,3$). Similarly, for each $H = \sum_{i=1}^3 z_i(a) H_i \in \mathfrak{h}_{\mathbb{Z}[a]} := \mathit{Span}_{\mathbb{Z}[a]}\big(H_1, H_2, H_3\big)$ — with $z_i(a) \in \mathbb{Z}[a]$ for all i — it is well-defined the element

$$h_H^{[a]}(t) \,:=\, t^H \,=\, \big(t^{H_1}\big)^{z_1(a)} \big(t^{H_2}\big)^{z_2(a)} \big(t^{H_3}\big)^{z_3(a)} \,\in\, \mathrm{GL}\big(V_{\Bbbk}(A)\big)\ , \quad \forall\ t \in P_a(A)$$

Definition 4.8. Let $H \in \mathfrak{h}_{\mathbb{Z}[a]}$ as above. We define the supergroup functor $h_H^{[a]}$ from (salg) to (groups) as

$$A \;\mapsto\; h_H^{[a]}(A) \;:=\; \big\{\, h_H^{[a]}(t) := t^H \;\big|\; t \in P(A_0) \big\}$$

By constrution, every functor $h_i^{[a]}$ is representable, namely by the Hopf algebra $\Bbbk\Big[\big\{d^{\pm a^k}\big\}_{k\in\mathbb{N}}\Big]$. Thus for each $H \in \mathfrak{h}_{\mathbb{Z}[a]}$ the supergroup functor $h_H^{[a]}$ is indeed an *affine supergroup* (actually, a classical affine group, indeed), which we shall mention as *multiplicative one-parameter (super)subgroup of a–type.*

4.5 Construction of Chevalley supergroups

In order to define our Chevalley supergroups, we first need the definition of a suitable *algebraic group* $\mathbf{G}_0$ associated to $\mathfrak{g} = D(2,1;a)$ and V.

First note that, by construction, the one-parameter supersubgroups $h_H(A)$ and $x_\alpha(A)$ (with $H \in \mathfrak{h}_{\mathbb{Z}}^0$, $\alpha \in \Delta_0$) altogether generate inside $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$ a *classical* affine groups, namely the (Chevalley-like) algebraic group associated to the semisimple Lie algebra $\mathfrak{g}_0$ — isomorphic to $\mathfrak{sl}(2)^{\oplus 3}$ — and its representation V. We denote it by $G_0'(A)$, so

$$G_0'(A) \;:=\; \Big\langle\, h_H(A)\,,\, x_\alpha(A) \;\Big|\; H \in \mathfrak{h}_{\mathbb{Z}}^0\,,\, \alpha\in\Delta_0 \Big\rangle$$

In particular, the h_H's generate $T'(A_0) := \big\langle\, h_H(A) \;\big|\, H \in \mathfrak{h}^0_{\mathbb{Z}} \,\big\rangle$, which is a maximal torus inside $G'_0(A)$. Both $A \mapsto G'_0(A)$ and $A \mapsto T'(A)$ are group-functors defined on (salg), which clearly factor through (alg) = $(\text{alg})_{\Bbbk}$, the category of unital commutative $\Bbbk$–algebras.

Note that G'_0 is a presheaf (cf. [4], Appendix A.5), hence we can define the functor $\mathbf{G}'_0$ as the sheafification of G'_0: on *local* algebras — in (alg) — the functor $\mathbf{G}'_0$ coincides with the functor of points of the Chevalley group-scheme associated with $\mathfrak{g}_0$ and V. Thus $\mathbf{G}'_0$ is just that functor of points, hence (in particular) it is representable.

Similarly, we can consider also the sheafification $\mathbf{T}'$ of T', yet $\mathbf{T}' = T'$.

Second, we consider the subgroup of $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$

$$T(A) \;:=\; \Big\langle\, \Big\{ h_H^{[a]}(A) \;\Big|\; H \in \mathfrak{h}_{\mathbb{Z}[a]} \Big\} \bigcup \, T'(A) \,\Big\rangle$$

which for various A in (salg) yields another functor $T : (\text{salg}) \longrightarrow (\text{groups})$. As T' and the various $h_H^{[a]}$'s are subgroups in $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$, we argue that T itself is a subgroup functor too.

We can now introduce the algebraic group $\mathbf{G}_0$ we were looking for:

Definition 4.9. For every $A \in (\text{salg})$, we let $G_0(A)$ be the subgroup

$$G_0(A) \;:=\; \Big\langle\, G'_0(A) \bigcup \, T(A) \,\Big\rangle$$

of $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$. We denote $G_0 : (\text{salg}) \longrightarrow (\text{groups})$ the supergroup functor which is the full subfunctor of $\mathrm{GL}\big(V_{\Bbbk}(A)\big)$ given on objects by $A \mapsto G_0(A)$. Also, we denote $\mathbf{G}_0 : (\text{salg}) \longrightarrow (\text{groups})$ the sheafification functor of G_0.

Note that both G_0 and $\mathbf{G}_0$ factor through (alg).

Proposition 4.10. *The supergroup functor* $\mathbf{G}_0$ *is representable, hence — as it factors through* (alg) *— it is an affine group.*

Proof. The groups $G'_0(A_0)$ and $T(A_0)$ are subgroups of $\mathrm{GL}(V_{\Bbbk})(A_0)$, and their mutual intersection is $G'_0(A_0) \cap T(A_0)$. Then the subgroup $G_0(A_0)$ is a fibered coproduct of $G'_0(A_0)$ and $T(A_0)$ over $T'(A_0)$. In down-to-earth terms, we can describe it as follows. Inside $\mathrm{GL}(V_{\Bbbk})(A_0)$, the subgroup $T(A_0)$ acts on $G'_0(A_0)$ by adjoint action, hence $G_0(A_0)$, generated by $T(A_0)$ and $G'_0(A_0)$, is a quotient of the semi-direct product $T(A_0) \ltimes G'_0(A_0)$. To be precise, we have the functorial isomorphism $G_0(A_0) \cong \big(T(A_0) \ltimes G_0'(A_0)\big)\Big/K(A_0)$ where

$$A \mapsto K(A_0) := \big\{ \big(k^{-1}, k\big) \,\big|\, t \in T(A_0) \cap G_0'(A_0) \big\}$$

defines — on (salg), through (alg) — a normal subgroup(-functor) of $T \ltimes G_0'$. So $G_0 \cong \big(T \ltimes G_0'\big)\Big/K$ as group-functors, hence $G_0 \cong \big(T \times G_0'\big)\Big/K$ as set-valued functors (i.e. forgetting the group structure) since $T \ltimes G_0' \cong T \times G_0'$ as set-valued functors. Taking sheafifications, as both $\mathbf{T}$ and $\mathbf{G}_0'$ are representable, we infer that the product $\mathbf{T} \times \mathbf{G}_0'$ is representable too. In addition, if $\mathbf{K}$ is the sheafification of the functor K, we see now that $\mathbf{K}$ as a subgroup of the algebraic group $\mathbf{T} \times \mathbf{G}_0'$ is *closed*.

To prove the last claim, let us revisit our construction. We started with a representation of $A \otimes_{\Bbbk} U_{\Bbbk}(\mathfrak{g})$ on $V_{\Bbbk}(A) := A \otimes_{\Bbbk} V_{\Bbbk}$ (cf. §4.1). By the very construction, the subgroup $\mathbf{G}_0'$ is nothing but the algebraic group associated to $\mathfrak{g}_0$ and V (as a $\mathfrak{g}_0$–module) by the classical Chevalley's construction: indeed, we have $\mathfrak{g}_0 \cong \mathfrak{sl}_2 \oplus \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$ where the three summands are given by $\mathfrak{sl}_2$–triples associated to positive even roots $2\,\varepsilon_i$ (cf. §2.3), and so $\mathbf{G}_0' \cong \mathbf{H}_1 \times \mathbf{H}_2 \times \mathbf{H}_3$ where $\mathbf{H}_i \in \big\{ \mathbf{SL}_2 \,, \mathbb{P}\mathbf{SL}_2 \big\}$ for each i. On the other hand, the subgroup $\mathbf{T}$ generated by the three one-parameter (super)subgroups of a–type $h^{[a]}_{H_i}$ — $i = 1, 2, 3$ — altogether is isomorphic to $P_a \times P_a \times P_a$. Recall also that the group P_a is represented by the $\Bbbk$–algebra $\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k \in \mathbb{N}}\big]$, while each $\mathbf{H}_i$ is represented by $\mathcal{O}(\mathbf{SL}_2) = \Bbbk\big[\mathrm{a}, \mathrm{b}, \mathrm{c}, \mathrm{d}\big]\Big/\big(\mathrm{ad} - \mathrm{bc} - 1\big)$ if $\mathbf{H}_i \cong \mathbf{SL}_2$, and by the unital subalgebra of $\mathcal{O}(\mathbf{SL}_2)$ generated by all products of any two elements in the set $\big\{\mathrm{a}, \mathrm{b}, \mathrm{c}, \mathrm{d}\big\}$ if $\mathbf{H}_i \cong \mathbb{P}\mathbf{SL}_2$.

Let's assume for a moment that $\mathbf{H}_i \cong \mathbf{SL}_2$ for all i. Then the identities

$$H_{2\,\varepsilon_1} = H_2 \;\; , \qquad H_{2\,\varepsilon_2} = 2H_1 - (1+a)H_2 - aH_3 \;\; , \qquad H_{2\,\varepsilon_3} = H_3$$

inside $\mathfrak{h}$ yield, for every $t_1, t_2, t_3 \in P_a(A)$, formal identities

$$h_{H_{2\varepsilon_1}}(t_1) = t_1^{H_{2\varepsilon_1}} = t_1^{H_2} = h^{[a]}_{H_2}(t_1) \;\; , \qquad h_{H_{2\varepsilon_3}}(t_3) = t_3^{H_{2\varepsilon_3}} = t_3^{H_3} = h^{[a]}_{H_3}(t_3)$$
$$h_{H_{2\varepsilon_2}}(t_2) = t_2^{2H_1 - (1+a)H_2 - aH_3} = h^{[a]}_{2H_1}(t_2) \cdot h^{[a]}_{-(1+a)H_2}(t_2) \cdot h^{[a]}_{-aH_3}(t_2)$$

Therefore, in the overall isomorphism $\mathbf{T} \times \mathbf{G}_0' \cong \big(P_a^{\times 3}\big) \times \big(\times_{i=1}^{3} \mathbf{H}_i\big)$, the subgroup $K(A_0)$ in $\mathbf{T} \times \mathbf{G}_0'$ — see above — corresponds in $\big(P_a^{\times 3}\big) \times \big(\times_{i=1}^{3} \mathbf{H}_i\big)$ to the subgroup

$$\left\{ \left(\big(t_2^{-2},\, t_1^{-1}\, t_2^{1+a},\, t_2^{a}\, t_3^{-1}\big) \,,\, \left(\begin{pmatrix} t_1 & 0 \\ 0 & t_1^{-1} \end{pmatrix}, \begin{pmatrix} t_2 & 0 \\ 0 & t_2^{-1} \end{pmatrix}, \begin{pmatrix} t_3 & 0 \\ 0 & t_3^{-1} \end{pmatrix} \right) \right) \,\middle|\, t \in P_a(A) \right\}$$

But this last subgroup is closed in $\big(P_a^{\times 3}\big) \times \big(\times_{i=1}^3 \mathbf{H}_i\big)$. Indeed, the latter is represented by the algebra $\mathcal{O}\big[\big(P_a^{\times 3}\big)\times\big(\times_{i=1}^3\mathbf{H}_i\big)\big] \cong \mathcal{O}\big[P_a\big]^{\otimes 3}\otimes\mathcal{O}\big[\mathbf{SL}_2\big]^{\otimes 3}$: to describe this algebra, let $\Bbbk\big[\big\{\ell_i^{\pm a^k}\big\}_{k\in\mathbb{N}}\big]$ be the i–th copy of $\Bbbk\big[\big\{\ell^{\pm a^k}\big\}_{k\in\mathbb{N}}\big] \cong \mathcal{O}\big[P_a\big]$ in this tensor product and let $\Bbbk\big[\mathrm{a}_j, \mathrm{b}_j, \mathrm{c}_j, \mathrm{d}_j\big]\Big/\big(\mathrm{a}_j\mathrm{d}_j - \mathrm{b}_j\mathrm{c}_j - 1\big) \cong \mathcal{O}\big[\mathbf{SL}_2\big]$ be the j–th copy of $\mathcal{O}\big[\mathbf{SL}_2\big]$. Then, by construction, the subgroup above is just the closed subscheme of $\big(P_a^{\times 3}\big) \times \big(\times_{i=1}^3 \mathbf{H}_i\big)$ defined by the ideal

$$I \;:=\; \Big(\, \ell_2\, \mathrm{a}_1 - 1 \;,\; \ell_1^2 \ell_2^{-1} \ell^{-a} \ell_3^{-a}\, \mathrm{a}_2 - 1 \;,\; \ell_3\, \mathrm{a}_3 - 1 \;,\; \mathrm{b}_1 \;,\; \mathrm{b}_2 \;,\; \mathrm{b}_3 \;,\; \mathrm{c}_1 \;,\; \mathrm{c}_2 \;,\; \mathrm{c}_3 \,\Big)$$

A similar analysis works too when some $\mathbf{H}_i$'s are isomorphic to $\mathbb{P}\mathbf{SL}_2$.

Finally, recall that $\mathbf{T} \times \mathbf{G}_0'$ is representable (hence it is an affine group scheme); then its quotients by closed normal subgroups are again affine groups. In particular, this holds for the quotient modulo $\mathbf{K}$ is representable (an affine group), hence also for the isomorphic functor $\mathbf{G}$. $\square$

We can now eventually define our Chevalley supergroups:

Definition 4.11. Let $\mathfrak{g}$ and V be as above. We call *Chevalley supergroup functor*, associated to $\mathfrak{g}$ and V , the functor $G : (\text{salg}) \longrightarrow (\text{grps})$ given by:

— if $A \in Ob\big((\text{salg})\big)$ we let $G(A)$ be the subgroup of $\mathrm{GL}\big(V_\Bbbk(A)\big)$ generated by $G_0(A)$ and the one-parameter subgroups $x_\beta(A)$ with $\beta \in \Delta_1$, that is

$$G(A) \;:=\; \Big\langle\, G_0(A)\,,\, x_\beta(A) \;\Big|\; \beta \in \Delta_1 \Big\rangle \;=\; \Big\langle T(A)\,,\, x_\delta(A) \;\Big|\; \delta \in \Delta \Big\rangle$$

where the second identity follows from the previous description of G_0 .

— if $\phi \in \mathrm{Hom}_{(\text{salg})}\big(A\,,B\big)$, then $\mathrm{End}_\Bbbk(\phi) : \mathrm{End}_\Bbbk\big(V_\Bbbk(A)\big) \longrightarrow \mathrm{End}_\Bbbk\big(V_\Bbbk(B)\big)$ (given on matrix entries by ϕ itself) respects the sum and the associative product of matrices. Then $\mathrm{End}_\Bbbk(\phi)$ clearly restricts to a group morphism $\mathrm{GL}\big(V_\Bbbk(A)\big) \longrightarrow \mathrm{GL}\big(V_\Bbbk(B)\big)$. The latter maps the generators of $G(A)$ to those of $G(B)$, hence restricts to a group morphism $G(\phi) : G(A) \longrightarrow G(B)$.

We call *Chevalley supergroup* — associated to $\mathfrak{g} = D(2,1;a)$ and V — the sheafification $\mathbf{G}$ of G (cf. [4], Appendix). Therefore $\mathbf{G} : (\text{salg}) \longrightarrow (\text{grps})$ is a sheaf functor such that $\mathbf{G}(A) = G(A)$ whenever $A \in (\text{salg})$ is *local*.

Finally, in order to stress the dependence on V, we shall sometimes also write G_V and $\mathbf{G}_V$ for G and $\mathbf{G}$ respectively.

Remarks 4.12.

(a) For $a \in \mathbb{Z}$, a construction of Chevalley supergroups of type $D(2,1;a)$ was given in [4] too, and coincides with the present one, because $P_a(A) = U(A_0)$ when $a \in \mathbb{Z}$ — for every $A \in (\text{salg})$.

(b) An alternative definition of Chevalley supergroups can be given by letting the subgroup (functor) $Up : A \mapsto Up(A_0)$ play the role of P_a, where $Up(A_0) := \big(1 + \mathfrak{N}(A_0)\big)$ is the subgroup of $U(A_0)$ of all *unipotent* elements of A_0 (with $\mathfrak{N}(A_0)$ being the nilradical of A_0). All our arguments and results from now on still stand valid as well. Nevertheless, choosing the option of the subgroup functor Up one does not recover the construction of [4] for $a \in \mathbb{Z}$, which instead is a "good feature" of the functor P_a, see *(a)* above.

4.6 Chevalley supergroups as affine supergroups

Our definition of the Chevalley supergroup $\mathbf{G}$ does not imply (at least not at first sight) that $\mathbf{G}$ is *representable*, in other words, that $\mathbf{G}$ is the functor of points of an affine supergroup scheme. In this section we prove this.

We begin with some more definitions:

Definition 4.13. For any $A \in (\text{salg})$, we define the subsets of $G(A)$

$$G_1(A) \;:=\; \Big\{ \textstyle\prod_{i=1}^{n} x_{\gamma_i}(\vartheta_i) \;\Big|\; n \in \mathbb{N} \,,\; \gamma_i \in \Delta_1 \,,\; \vartheta_i \in A_1 \Big\}$$

$$G_0^{\pm}(A) \;:=\; \Big\{ \textstyle\prod_{i=1}^{n} x_{\alpha_i}(t_i) \;\Big|\; n \in \mathbb{N} \,,\; \alpha_i \in \Delta_0^{\pm} \,,\; t_i \in A_0 \Big\}$$

$$G_1^{\pm}(A) \;:=\; \Big\{ \textstyle\prod_{i=1}^{n} x_{\gamma_i}(\vartheta_i) \;\Big|\; n \in \mathbb{N} \,,\; \gamma_i \in \Delta_1^{\pm} \,,\; \vartheta_i \in A_1 \Big\}$$

$$G^{\pm}(A) \;:=\; \left\{ \prod_{i=1}^{n} x_{\beta_i}(\mathbf{t}_i) \;\middle|\; n \in \mathbb{N} \,,\; \beta_i \in \Delta^{\pm},\; \mathbf{t}_i \in A_0 \cup A_1 \right\} \;=\; \big\langle G_0^{\pm}(A) \,, G_1^{\pm}(A) \big\rangle$$

Moreover, fixing any total order $\preceq$ on $\Delta_1^{\pm}$, and letting $N_{\pm} = \big|\Delta_1^{\pm}\big|$, we set

$$G_1^{\pm,<}(A) \;:=\; \Big\{ \textstyle\prod_{i=1}^{N_{\pm}} x_{\gamma_i}(\vartheta_i) \;\Big|\; \gamma_1 \prec \cdots \prec \gamma_{N_{\pm}} \in \Delta_1^{\pm} \,,\; \vartheta_1, \dots, \vartheta_{N_{\pm}} \in A_1 \Big\}$$

and for any total order $\preceq$ on Δ_1, and letting $N := \big|\Delta\big| = N_+ + N_-$, we set

$$G_1^{<}(A) \;:=\; \Big\{ \textstyle\prod_{i=1}^{N} x_{\gamma_i}(\vartheta_i) \;\Big|\; \gamma_1 \prec \cdots \prec \gamma_N \in \Delta_1 \,,\; \vartheta_1, \dots, \vartheta_N \in A_1 \Big\}$$

By the way, note that $N_\pm = 4$ and $N = 8$ for $\mathfrak{g} = D(2,1;a)$.

Similar notations will denote the sheafifications $\mathbf{G}_1$, $\mathbf{G}^\pm$, $\mathbf{G}_0^\pm$, $\mathbf{G}_1^\pm$, etc.

We shall start by studying the commutation relations of the generators of Chevalley groups. As a matter of notation, when $\varGamma$ is any group and $g, h \in \varGamma$ we denote by $(g,h) := g\,h\,g^{-1}\,h^{-1}$ their commutator in $\varGamma$.

Lemma 4.14.

(a) Let $\alpha \in \Delta_0$, $\gamma \in \Delta_1$, $A \in (\mathrm{salg})$ *and* $t \in A_0$, $\vartheta \in A_1$. *Then there exist* $c_s \in \mathbb{Z}$ *such that*

$$\big(x_\gamma(\vartheta) \, , \, x_\alpha(t) \big) \; = \; \prod\nolimits_{s>0} x_{\gamma+s\,\alpha}\big(c_s \, t^s \vartheta \big) \;\; \in \;\; G_1(A)$$

(the product being finite). Indeed, with $\varepsilon_k = \pm 1$ *and* r *as in §3.4, part (2),*

$$\big(1 + \vartheta \, X_\gamma \, , \, x_\alpha(t) \big) \; = \; \prod\nolimits_{s>0} \Big(1 + \prod\nolimits_{k=1}^{s} \varepsilon_k \cdot \tbinom{s+r}{r} \cdot t^s \vartheta \, X_{\gamma+s\,\alpha} \Big)$$

where the factors in the product are taken in any order (as they do commute).

(b) Let $\gamma, \delta \in \Delta_1$, $A \in (\mathrm{salg})$, $\vartheta, \eta \in A_1$. *Then (notation of Definition 3.1)*

$$\big(x_\gamma(\vartheta) \, , \, x_\delta(\eta) \big) \;\; = \;\; x_{\gamma+\delta}\big(-c_{\gamma,\delta} \, \vartheta \, \eta \big) \;\; = \;\; \big(1 - c_{\gamma,\delta} \, \vartheta \, \eta \, X_{\gamma+\delta} \big) \;\; \in \;\; G_0(A)$$

if $\delta \neq -\gamma$ *; otherwise, for* $\delta = -\gamma$, *we have*

$$\big(x_\gamma(\vartheta) \, , \, x_{-\gamma}(\eta) \big) \;\; = \;\; \big(1 - \vartheta \, \eta \, H_\gamma \big) \;\; = \;\; h_{H_\gamma}^{[a]}\big(1 - \vartheta \, \eta \big) \;\; \in \;\; G_0(A)$$

(c) Let $\beta \in \Delta$, $A \in (\mathrm{salg})$, $\mathbf{u} \in A_0 \cup A_1$. *Then*

$$h_{H_0}(t) \; x_\beta(\mathbf{u}) \; h_{H_0}(t)^{-1} = \; x_\beta\big(t^{\beta(H_0)} \, \mathbf{u} \big) \;\in\; G_{p(\beta)}(A) \quad \forall \;\; H_0 \in \mathfrak{h}_{\mathbb{Z}}^0 \, , \, t \in U(A_0)$$

$$h_H^{[a]}(t) \; x_\beta(\mathbf{u}) \; h_H^{[a]}(t)^{-1} \; = \; x_\beta\big(t^{\beta(H)} \, \mathbf{u} \big) \;\in\; G_{p(\beta)}(A) \quad \forall \;\; H \in \mathfrak{h}_{\mathbb{Z}[a]} \, , \, t \in P_a(A_0)$$

where $p(\beta) := s$, *by definition, if and only if* $\beta \in \Delta_s$.

Proof. Like for [4], Lemma 5.11, these results follow directly from the classical ones in [11], pg. 22 and 29, and simple calculations, using the relations in §3.4 and the identity $(\vartheta\eta)^2 = -\vartheta^2\,\eta^2 = 0$. In addition, for *(b)* in the present case we have also to take into account that $(1 - \vartheta\,\eta) \in \big(1 + \mathfrak{N}(A_0) \big) \subseteq P_a(A)$ for all $\vartheta, \eta \in A_1$, so $h_{H_\gamma}^{[a]}\big(1 - \vartheta\,\eta\big)$ is well-defined and equal to $\big(1 - \vartheta\,\eta\,H_\gamma \big)$. □

Now, with our definition of Chevalley groups at hand and the commutation rules among their generators available – as given in Lemma 4.14 above — one can reproduce whatever was done in [4], §5.3. The results then will be the same, and, for their proofs, exactly the same arguments will work, just adapting them to the present context. Just some minimal changes are due to a couple of facts: first, the presence among generators of the multiplicative one-parameter of type a, which are handled like the classical ones using Lemma 4.14; second, several possible shortcuts and simplifications (compared to [4]) are possible, as the structure of $\mathfrak{g} = D(2,1;a)$ is much simpler than that of the general classical Lie superalgebra. Therefore, in the following I shall bound myself to list the results we get (essentially, the main steps in the line of arguing of [4]), as the proof can be easily recovered from [4], §3.

Our first result is:

Theorem 4.15. *For any* $A \in (\mathrm{salg})$ *, there exist set-theoretic factorizations*

$$\begin{array}{llll} \textit{(a)} & G(A) \, = \, G_0(A)\, G_1(A) \quad , & & G(A) \, = \, G_1(A)\, G_0(A) \\ \textit{(b)} & G^\pm(A) \, = \, G_0^\pm(A)\, G_1^\pm(A) \quad , & & G^\pm(A) \, = \, G_1^\pm(A)\, G_0^\pm(A) \\ \textit{(c)} & G(A) \, = \, G_0(A)\, G_1^<(A) \quad , & & G(A) \, = \, G_1^<(A)\, G_0(A) \end{array}$$

Claim *(a)* above is a group-theoretical counterpart for G of the splitting $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ — i.e. a super-analogue of the Cartan decomposition for reductive groups — while *(b)* is a similar result for G^+ and G^-. Claim *(c)* improves *(a)*: like in [4] (Theorem 4.13), it follows from Theorem 4.15 just reordering the factors in $G_1(A)$ by means of the commutation rules in Lemma 4.14.

With a more careful analysis, one then improves the previous results to get the following, key one (the analogue of Corollary 5.18 in [4]):

Theorem 4.16. *The group product yields functor isomorphisms*

$$G_0 \times G_1^{-,<} \times G_1^{+,<} \xrightarrow{\cong} G \quad , \qquad \mathbf{G}_0 \times \mathbf{G}_1^{-,<} \times \mathbf{G}_1^{+,<} \xrightarrow{\cong} \mathbf{G}$$

as well as those obtained by permuting the $(-)$*-factor and the* $(+)$*-factor and/or moving the* (0)*-factor to the right. Moreover, all these induce similar functor isomorphisms with the left-hand side obtained by permuting the factors above, like* $G_1^{+,<} \times G_0 \times G_1^{-,<} \xrightarrow{\cong} G$ *,* $\mathbf{G}_1^{-,<} \times \mathbf{G}_0 \times \mathbf{G}_1^{+,<} \xrightarrow{\cong} G$ *, etc.*

Notes that the isomorphisms in Theorem 4.16 can be thought of as super-analogue of the classical *big cell decomposition* for reductive algebraic groups.

The same technique used to prove Theorem 4.16 also yields the following:

Proposition 4.17. *The functors* $G_1^{\pm,<} : (\mathrm{salg}) \longrightarrow (\mathrm{sets})$ *are representable: they are the functor of points of the superscheme* $\mathbb{A}_{\Bbbk}^{0|N_\pm}$*, with* $N_\pm = \left|\Delta_1^\pm\right|$ *. In particular they are sheaves, hence* $G_1^{\pm,<} = \mathbf{G}_1^{\pm,<}$ *.*

Indeed, this holds since natural transformations $\Psi^\pm : \mathbb{A}_{\Bbbk}^{0|N_\pm} \longrightarrow G_1^{\pm,<}$ exist, by definition, given on objects by $\Psi^\pm(A) : \mathbb{A}_{\Bbbk}^{0|N_\pm}(A) \longrightarrow G_1^{\pm,<}(A)$, $(\vartheta_1, \dots, \vartheta_{N_\pm}) \mapsto \prod_{i=1}^{N_\pm} x_{\gamma_i}(t_i)$ (and obvious on morphisms). One then proves that these $\Psi^\pm$ are isomorphisms of functors.

Finally, we can prove that the Chevalley supergroups are affine:

Theorem 4.18. *Every Chevalley supergroup* $\mathbf{G}$ *is an affine supergroup.*

Indeed, this is a direct consequence of the last two results, as they imply that the group-functor $\mathbf{G}$ is (isomorphic to) the direct product of three representable group-functors, hence in turn it is representable as well, which entails that it is an affine supergroup.

Another immediate consequence (of the same results) is the following, which improves, for Chevalley supergroups, a more general result proved by Masuoka (cf. [9], Theorem 4.5) in the algebraic-supergeometry setting (see also [13], and references therein, for the complex-analytic case).

Proposition 4.19. *For any Chevalley supergroup* $\mathbf{G}$*, there are isomorphisms of commutative superalgebras*

$$\begin{aligned} \mathcal{O}(\mathbf{G}) \;&\cong\; \mathcal{O}(\mathbf{G}_0) \otimes \mathcal{O}\big(\mathbf{G}_1^{-,<}\big) \otimes \mathcal{O}\big(\mathbf{G}_1^{+,<}\big) \;\cong \\ &\cong\; \mathcal{O}(\mathbf{G}_0) \otimes \Bbbk\big[\xi_1, \dots, \xi_{N_-}\big] \otimes \Bbbk\big[\chi_1, \dots, \chi_{N+}\big] \end{aligned}$$

where $N_\pm = \left|\Delta_1^\pm\right|$*, the subalgebra* $\mathcal{O}(\mathbf{G}_0)$ *is totally even, and* $\xi_1, \dots, \xi_{N_-}$ *and* $\chi_1, \dots, \chi_{N_+}$ *are odd elements.*

Finally, one has the following result for any $A \in (\mathrm{salg})$ which is *the central extension of the commutative algebra* A_0 *by the* A_0*–module* A_1 :

Proposition 4.20. *Let G be a Chevalley supergroup functor, and let $\mathbf{G}$ be its associated Chevalley supergroup. Assume $A \in (\mathrm{salg})$ is such that $A_1^2 = \{0\}$.*

Then $G_1^+(A)$, $G_1^-(A)$ and $G_1(A)$ are normal subgroups of $G(A)$, with

$$\begin{gathered} G_1^{\pm}(A) \; = \; G_1^{\pm,<}(A) \; \cong \; \mathbb{A}_{\Bbbk}^{0|N_\pm}(A) \qquad\qquad \textit{with } \; N_\pm = \big|\Delta_1^\pm\big| \\ G_1(A) \; = \; G_1^-(A) \cdot G_1^+(A) \; = \; G_1^+(A) \cdot G_1^-(A) \\ G_1(A) \; \cong \; G_1^-(A) \times G_1^+(A) \; \cong \; G_1^+(A) \times G_1^-(A) \; \cong \; \mathbb{A}_{\Bbbk}^{0|N_-}(A) \times \mathbb{A}_{\Bbbk}^{0|N_+}(A) \end{gathered}$$

(where "$\cong$" means isomorphic as groups), the group structure on $\mathbb{A}_{\Bbbk}^{0|N_\pm}(A)$ being the obvious one. In particular, $G(A)$ is the semidirect product, namely $G(A) \cong G_0(A) \ltimes G_1(A) \cong G_0(A_0) \ltimes \Big(\mathbb{A}_{\Bbbk}^{0|N_-}(A) \times \mathbb{A}_{\Bbbk}^{0|N_+}(A)\Big)$, of the classical group $G_0(A_0)$ with the totally odd affine superspace $\mathbb{A}_{\Bbbk}^{0|N_-}(A) \times \mathbb{A}_{\Bbbk}^{0|N_+}(A)$.

Similar results hold with a symbol "$\mathbf{G}$" replacing "G" everywhere.

5 Independence of the construction and Lie's Third Theorem

In this section, we discuss the dependence of Chevalley supergroups $\mathbf{G}_V$ on the $\mathfrak{g}$–modules V, and a super-analogue of Lie's Third Theorem for $\mathbf{G}_V$ and its converse. Once more, we shall essentially repeat the statements of [4], §§5.4–5 (the proof being the same).

5.1 Independence of Chevalley and Kostant superalgebras

The construction of Chevalley groups depend on the finite dimensional, rational $\mathfrak{g}$–representation V fixed from scratch, and on an admissible $\mathbb{Z}$–lattice M in V. I show now how it depends on V: as a consequence, one finds that it is in fact *independent of M*. Once again, I stick to statements, as the proofs follow the same arguments as in [4], §§5.4–5.

Let $\mathbf{G}'$ and $\mathbf{G}$ be two Chevalley supergroups obtained by the same $\mathfrak{g}$, possibly with a different choice of the representation. We denote with X_α and with X'_α respectively the elements of the Chevalley basis in $\mathfrak{g}$ identified (as usual) with their images under the two representations of $\mathfrak{g}$.

Our first result is technical, yet important:

Lemma 5.1. *Let* $\phi : \mathbf{G} \longrightarrow \mathbf{G}'$ *be a morphism of Chevalley supergroups such that for all local superalgebras* A *we have* $\phi_A\big(\mathbf{G}_0(A)\big) = \mathbf{G}'_0(A)$ *and* $\phi_A\big(1 + \vartheta\, X_\beta\big) = 1 + \vartheta\, X'_\beta$ *for all* $\beta \in \Delta_1$, $\vartheta \in A_1$. *Then* $Ker(\phi_A) \subseteq \mathbf{T}$, *where* $\mathbf{T}$ *is the maximal torus in the algebraic group* $\mathbf{G}_0 \subseteq \mathbf{G}$ *(see §4.5).*

Then let L_V be the lattice spanned by the weights (of $\mathfrak{h}$) in the $\mathfrak{g}$–representation V. Then the relation between Chevalley supergroups corresponding to different weight lattices is the same as in the classical setting:

Theorem 5.2. *Let* $\mathbf{G}$ *and* $\mathbf{G}'$ *be two Chevalley supergroups constructed using two representations* V *and* V' *of the same* $\mathfrak{g}$ *over the same field* $\mathbb{K}$ *(as in §4.1). If* $L_V \supseteq L_{V'}$, *then there exists a unique morphism* $\phi : \mathbf{G} \longrightarrow \mathbf{G}'$ *such that* $\phi_A\big(1{+}\,\vartheta\, X_\alpha\big) = 1 + \vartheta\, X'_\alpha$, *and* $Ker(\phi_A) \subseteq Z\big(\mathbf{G}(A)\big)$, *for every local algebra* A . *Moreover,* ϕ *is an isomorphism if and only if* $L_V = L_{V'}$.

As a direct consequence, we have the following "independence result":

Corollary 5.3. *Every Chevalley supergroup* $\mathbf{G}_V$ *is independent — up to isomorphism — of the choice of an admissible lattice* M *of* V *considered in the very construction of* $\mathbf{G}_V$ *itself.*

5.2 Lie's Third Theorem

In the present context, the analogue of "Lie's Third Theorem" concerns the question of whether the tangent Lie superalgebra of our supergroups $\mathbf{G}$ is $\mathfrak{g} = D(2,1;a)$. We shall now cope with this problem.

Let now $\Bbbk$ be a *field*, with $\mathrm{char}(\Bbbk) \neq 2,3$. Let $\mathbf{G}_V$ be a Chevalley supergroup scheme over $\Bbbk$, built out of a $\mathfrak{g} = D(2,1;a)$ and a rational $\mathfrak{g}$–module V as in §4.5. In §4.1, we have constructed the Lie superalgebra $\mathfrak{g}_\Bbbk := \Bbbk \otimes_{\mathbb{Z}_a} \mathfrak{g}_V$ over $\Bbbk$ starting from the $\mathbb{Z}_a$–lattice $\mathfrak{g}_V$. We now show that the affine supergroup $\mathbf{G}_V$ has $\mathfrak{g}_\Bbbk$ as its tangent Lie superalgebra.

In [4], §5.5, one can read how to associate a Lie superalgebra to a supergroup scheme (in a functorial way) — see also [3] for more details.

Let $A \in (\mathrm{salg})$ and let $A[\epsilon] := A[x]\big/\big(x^2\big)$ be the *super*algebra of dual numbers, in which $\epsilon := x \mod \big(x^2\big)$ is taken to be *even*. We have that

$A[\epsilon] = A \oplus \epsilon A$, and there are two natural morphisms $i : A \longrightarrow A[\epsilon]$, $a \overset{i}{\mapsto} a$, and $p : A[\epsilon] \longrightarrow A$, $(a + \epsilon\, a') \overset{p}{\mapsto} a$, such that $p \circ i = \mathrm{id}_A$.

Definition 5.4. For each supergroup scheme G, consider the homomorphism $G(p) : G(A(\epsilon)) \longrightarrow G(A)$. Then there is a supergroup functor

$$\mathrm{Lie}(G) : (\mathrm{salg}) \longrightarrow (\mathrm{sets}) , \qquad \mathrm{Lie}(G)(A) := \mathit{Ker}(G(p))$$

One shows that the functor Lie(G) is representable, and can be identified with (the functor of points of) the tangent space at the identity of G, namely $\mathit{Lie}(G)(A) = (A \otimes T_{1_G})_0$, where $T_{1_G} := \mathfrak{m}_{G,1_G} \big/ \mathfrak{m}^2_{G,1_G}$ is a super vector space, $\mathfrak{m}_{G,1_G}$ being the maximal ideal of the local algebra $\mathcal{O}_{G,1_G}$. Then by an abuse of notation one uses the same symbol Lie(G) to denote both the functor and the underlying super vector space representing it.

One also proves that Lie(G) has a Lie superalgebra structure: in other terms, the functor $\mathrm{Lie}(G) : (\mathrm{salg}) \longrightarrow (\mathrm{sets})$ is Lie algebra valued.

Now we compare with $\mathfrak{g} = D(2,1;a)$ the Lie superalgebra $\mathrm{Lie}(\mathbf{G}_V)$ of any of our Chevalley supergroups $\mathbf{G}_V$: the outcome is (cf. [4], Theorem 5.32):

Theorem 5.5. *If $\mathbf{G}_V$ is a Chevalley supergroup built upon $\mathfrak{g}$ and V, then* $\mathrm{Lie}(\mathbf{G}_V) = \mathfrak{g}$ *as functors with values in* (Lie-alg).

Remark 5.6. In the proof of Theorem 5.5 above, one uses also the (trivial to prove) fact that the (classical) affine group $P_a : (\mathrm{alg}) \longrightarrow (\mathrm{groups})$ has tangent Lie algebra the functor $\mathrm{Lie}(P_a) : (\mathrm{alg}) \longrightarrow (\mathrm{Lie\text{-}alg})$, $A \mapsto A$. The same also holds for the affine group $Up : (\mathrm{alg}) \longrightarrow (\mathrm{groups})$, which is relevant for Remark 4.12*(b)*.

References

[1] J. Brundan, A. Kleshchev, *Modular representations of the supergroup Q(n), I*, J. Algebra **206** (2003), 64–98.

[2] J. Brundan, J. Kujava, *A New Proof of the Mullineux Conjecture*, J. Alg. Combinatorics **18** (2003), 13–39.

[3] L. Caston, R. Fioresi, *Mathematical Foundation of Supersymmetry*, preprint `arXiv:0710.5742v1` [math.RA] (2007).

[4] R. Fioresi, F. Gavarini, *Chevalley Supergroups*, Trans. Amer. Math. Soc. (to appear) — preprint `arXiv:0808.0785` [math.RA] (2008).

[5] L. Frappat, P. Sorba, A. Sciarrino, *Dictionary on Lie algebras and superalgebras*, Academic Press, Inc., San Diego, CA, 2000.

[6] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Graduate Texts in Math., 9, Springer-Verlag, New York, 1972.

[7] K. Iohara, Y. Koga, *Central extensions of Lie Superalgebras*, Comment. Math. Helv. **76** (2001), 110–154.

[8] V. G. Kac, *Lie superalgebras*, Adv. in Math. **26** (1977), 8–26.

[9] Masuoka, A. *The fundamental correspondences in super affine groups and super formal groups.* J. Pure Appl. Algebra 202, No.1-3, 284-312, 2005.

[10] M. Scheunert, *The Theory of Lie Superalgebras*, Lecture Notes in Math. **716**, Springer-Verlag, Berlin-Heidelberg-New York, 1979.

[11] R. Steinberg, *Lectures on Chevalley groups*, Yale University, New Haven, Conn., 1968.

[12] B. Shu, W. Wang, *Modular representations of the ortho-symplectic supergroups*, Proc. Lond. Math. Soc. (3) **96** (2008), no. 1, 251–271.

[13] E. G. Vishnyakova, *On complex Lie supergroups and homogeneous split supermanifolds*, preprint `arXiv:0908.1164v1` [math.DG] (2009).

[14] V. S. Varadarajan, *Supersymmetry for mathematicians: an introduction*, Courant Lecture Notes **1**, AMS, 2004.